
Accepted Manuscript

Title: Determination and prediction of solubilities of active pharmaceutical ingredients in selected organic solvents

Author: Hiroyuki Matsuda Kanako Mori Mariko Tomioka Nozomi Kariyasu Toshiro Fukami Kiyofumi Kurihara Katsumi Tochigi Kazuo Tomono



PII:	S0378-3812(15)30047-9
DOI:	http://dx.doi.org/doi:10.1016/j.fluid.2015.07.032
Reference:	FLUID 10684
To appear in:	Fluid Phase Equilibria
Received date:	9-4-2015
Revised date:	13-7-2015
Accepted date:	15-7-2015

Please cite this article as: Hiroyuki Matsuda, Kanako Mori, Mariko Tomioka, Nozomi Kariyasu, Toshiro Fukami, Kiyofumi Kurihara, Katsumi Tochigi, Kazuo Tomono, Determination and prediction of solubilities of active pharmaceutical ingredients in selected organic solvents, *Fluid Phase Equilibria* (2015), http://dx.doi.org/10.1016/j.fluid.2015.07.032

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

- > This study investigates measurements of solubilities of APIs.
- > Solubilities of APIs in different organic solvents were measured.
- > Solubilities of famotidine with two polymorphs were also investigated.
- > Experimental solubility data were represented by NRTL and UNIQUAC models.
- > Solubilities of salicylic acid were predicted by Pharma Modified UNIFAC.

>ر

Download English Version:

https://daneshyari.com/en/article/6619645

Download Persian Version:

https://daneshyari.com/article/6619645

Daneshyari.com