A sandwich-type electrochemical aptasensor for determination of MUC 1 tumor marker based on PSMA-capped PFBT dots platform and high conductive rGO-N′1,N′3-dihydroxymalonimidamide/thionine nanocomposite as a signal tag

Leila Farzin, Sodeh Sadjadi, Mojtaba Shamsipur, Ammar Chabok, Shahab Sheibani

PII: S1572-6657(17)30789-0
DOI: doi:10.1016/j.jelechem.2017.11.011
Reference: JEAC 3641

To appear in: *Journal of Electroanalytical Chemistry*

Received date: 5 October 2017
Revised date: 31 October 2017
Accepted date: 5 November 2017

Please cite this article as: Leila Farzin, Sodeh Sadjadi, Mojtaba Shamsipur, Ammar Chabok, Shahab Sheibani , A sandwich-type electrochemical aptasensor for determination of MUC 1 tumor marker based on PSMA-capped PFBT dots platform and high conductive rGO-N′1,N′3-dihydroxymalonimidamide/thionine nanocomposite as a signal tag. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Jeac(2017), doi:10.1016/j.jelechem.2017.11.011

A sandwich-type electrochemical aptasensor for determination of MUC 1 tumor marker based on PSMA-capped PFBT dots platform and high conductive rGO-N'^{1},N'^{3}-dihydroxymalonimidamide/thionine nanocomposite as a signal tag

Leila Farzin*[a], Sodeh Sadjadi[b], Mojtaba Shamsipur[c], Ammar Chabok[d], Shahab Sheibani[a]

[a]*Radiation Application Research School, Nuclear Science and Technology Research Institute, Tehran, Iran*
[b]*Material and Nuclear fuel cycle school, Nuclear Science and Technology Research Institute, Tehran, Iran*
[c]*Department of Chemistry, Razi University, Kermanshah, Iran*
[d]*Department of Chemistry, Tarbiat Modares University, Tehran, Iran*