Accepted Manuscript

Title: Rapid reduction<!--<query id="Q1">Please check the document heading for correctness.</query>--> of dye pollutants and hexavalent chromium by silver-sulphur oxido-vanadium cluster

Authors: Biraj Das, Mukesh Sharma, Jugal C. Sarmah, Kusum

K. Bania

PII: S2213-3437(17)30404-9

DOI: http://dx.doi.org/10.1016/j.jece.2017.08.020


Reference: JECE 1815

To appear in:

Received date: 12-3-2017 Revised date: 12-8-2017 Accepted date: 15-8-2017

Please cite this article as: Biraj Das, Mukesh Sharma, Jugal C.Sarmah, Kusum K.Bania, Rapid reduction of dye pollutants and hexavalent chromium by silver-sulphur oxido-vanadium cluster, Journal of Environmental Chemical Engineeringhttp://dx.doi.org/10.1016/j.jece.2017.08.020

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ACCEPTED MANUSCRIPT


Rapid Reduction of Dye Pollutants and Hexavalent Chromium by Silver-Sulphur Oxido-Vanadium Cluster

Biraj Das, a Mukesh Sharma, b Jugal C. Sarmah, Kusum K. Bania*

Department of Chemical Sciences, Tezpur University, Napaam, Tezpur, Assam, India, 784028

Contact Author

Email: kusum@tezu.ernet.in Mob. No: 09859929360



Reduction of pollutant dye and Cr(VI) was achieved within few second with AgVT catalyst

Abstract: Very fast reduction of methylene blue, methyl orange and chromium (VI) was achieved with silver-sulphur-oxido-vanadium cluster in presence of sodium borohydride as reducing agent. With 10 mg of the catalyst and 100µL of 3 mM solution of sodium borohydride, methylene blue solution of 0.01mM concentration could be reduced within 4-5 sec. While with 10 mg of catalyst and 20 mL of sodium borohydride, 30 mg of methylene blue dissolved in water got decolourized within 10 min. Aerial oxidation of reduced leuco methylene blue species to methylene blue was seen after 3h on standing the reduced colourless solution of methylene blue. Prominent reduction of either methylene blue or

Download English Version:

https://daneshyari.com/en/article/6664239

Download Persian Version:

https://daneshyari.com/article/6664239

<u>Daneshyari.com</u>