

Accepted Manuscript

High pressure processing of barramundi (*Lates calcarifer*) muscle before freezing: the effects on selected physicochemical properties during frozen storage

Binh Q. Truong, Roman Buckow, Minh H. Nguyen, Costas E. Stathopoulos



PII: S0260-8774(15)00373-8

DOI: [10.1016/j.jfoodeng.2015.08.020](https://doi.org/10.1016/j.jfoodeng.2015.08.020)

Reference: JFOE 8297

To appear in: *Journal of Food Engineering*

Received Date: 9 February 2015

Revised Date: 11 August 2015

Accepted Date: 12 August 2015

Please cite this article as: Truong, B.Q., Buckow, R., Nguyen, M.H., Stathopoulos, C.E., High pressure processing of barramundi (*Lates calcarifer*) muscle before freezing: the effects on selected physicochemical properties during frozen storage, *Journal of Food Engineering* (2015), doi: 10.1016/j.jfoodeng.2015.08.020.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

High pressure processing of barramundi (*Lates calcarifer*) muscle before freezing: the effects on selected physicochemical properties during frozen storage

Binh Q. Truong^{1,4}, Roman Buckow², Minh H. Nguyen^{1,3*}, Costas E. Stathopoulos^{1,5}

¹ School of Environmental and Life Sciences, The University of Newcastle, Australia

² Commonwealth Scientific and Industrial Research Organisation, Food and Nutrition Flagship, Australia

³ School of Science and Health, University of Western Sydney, Australia

⁴ Faculty of Fisheries, Nong Lam University, Vietnam

⁵ Current address: Faculty of Bioscience Engineering, Ghent University Global Campus, Incheon, South Korea

*Corresponding Author:

Minh H. Nguyen

School of Environmental and Life Sciences, the University of Newcastle, Australia

Email: minh.nguyen@newcastle.edu.au

Highlights:

- Treatment at 150-200 MPa for 3 min before freezing improves the quality of frozen barramundi muscle.
- Treatment ≥ 250 MPa for 3 min resulted in cooked appearance and increased hardness of barramundi muscle.

Download English Version:

<https://daneshyari.com/en/article/6664945>

Download Persian Version:

<https://daneshyari.com/article/6664945>

[Daneshyari.com](https://daneshyari.com)