

Bacterial infection and pre-treatment with 24-epibrassinolide markedly affect the heat emission and membrane permeability of rape cotyledons

Anna Janeczko^{a,*}, István Tóbiás^b, Andrzej Skoczowski^a,
Franciszek Dubert^a, Gábor Gullner^b, Balázs Barna^b

^a Department of Biology of Flowering, The Franciszek Górski Institute of Plant Physiology,
Polish Academy of Sciences, 21 Niezapominajek St., 30-239 Kraków, Poland

^b Plant Protection Institute, Hungarian Academy of Sciences, 15 Herman Otto St., 1525 Budapest, Hungary

Available online 19 January 2007

Abstract

Changes in metabolic activity in spring oilseed rape (*Brassica napus* L.) seedlings during bacterial infection were determined by measuring heat emission with an isothermal calorimeter. Heat emission was markedly increased in cotyledons infected with the incompatible bacterium *Pseudomonas syringae* pv. *syringae*. One day after inoculation, visible signs of necrosis were detected on the infected cotyledons. This indicates that the hypersensitive response had been activated. Heat emission continued to increase as the hypersensitive response progressed, reflecting an increase in the metabolic rate in the infected tissue. With both of the cultivars tested heat emission by infected cotyledons was increased by pre-treatment with BR₂₇. Membrane permeability as measured by ion leakage also rapidly increased during the first 3 days of infection. In uninfected cotyledons pre-treated with BR₂₇, membrane permeability was slightly increased. In infected cotyledons not pre-treated with BR₂₇, both visible necrosis and membrane permeability rapidly increased during the first 2 or 3 days after inoculation. In infected cotyledons pre-treated with BR₂₇, membrane permeability was significantly reduced by the second or third day after inoculation. This confirms that BR₂₇ plays a role in preventing damage to plant cell membranes caused by bacterial infection during the hypersensitive response.

© 2007 Elsevier B.V. All rights reserved.

Keywords: Bacterial infection; *Brassica napus* L.; Isothermal calorimetry; Stress resistance; 24-Epibrassinolide

1. Introduction

Brassinosteroids are steroidal plant hormones that stimulate plant growth and protect plants against various stressors, including heavy metals, salt, heat shock, cold and pathogens [1–4]. Although they were discovered in the 1970s, little is known about the role brassinosteroids play in combating microbial infections in plants. The hypersensitive response is one of the best characterized defense mechanisms in infected host plants. The hypersensitive response involves programmed cell death, which causes rapid necrosis in infected tissues [5]. During the hypersensitive response, cell permeability rapidly increases as the cell membranes break down. Changes in metabolism also take place in plants during microbial infection. Calorimetric measurement of heat emission has been used to study metabolic changes in plants responding to various other stressors, includ-

ing salt, drought and herbicides [6–9]. The aim of this study was to determine the effects of 24-epibrassinolide (BR₂₇) on heat emission and cell permeability in oilseed rape seedlings artificially infected with the incompatible pathogenic bacterium *Pseudomonas syringae* pv. *syringae*.

2. Material and methods

2.1. Plant material and pre-treatment with BR₂₇

The study was carried out using two cultivars of spring oilseed rape (*Brassica napus* L.): ‘Licosmos’ and ‘Huzar’. After sowing, the plants were kept in a greenhouse for 14 days, by which time both cotyledons and the first leaf had developed. The cotyledons were then painted with a solution containing 100 nM BR₂₇, prepared from a stock solution containing 2 mM BR₂₇ in 96% ethanol. The BR₂₇ used in this study was purchased from Sigma (Poznań, Poland). The cotyledons of the control plants were painted with distilled water. A suspension of *P. syringae* pv. *syringae* containing 10⁸ cfu cm^{−3} was then injected into the

* Corresponding author. Tel.: +48 12 425 33 01/18 33; fax: +48 12 425 18 44.
E-mail address: ania@belanna.strefa.pl (A. Janeczko).

cotyledons by applying gentle pressure with the help of a plastic syringe not fitted with a needle. The concentration of the bacterial suspension was determined with the help of a spectrophotometer.

2.2. Measurement of heat emission

Rate of heat flow from the cotyledons was measured with the help of an isothermal LKB-2277 Bioactivity Monitor (Thermo-metric, Järfälla, Sweden). One day after the plants were treated with BR₂₇ and inoculated with bacterial suspension, the cotyledons were cut, placed in an ampule, and allowed to equilibrate for 20 min before analysis. Measurements were carried out for 10 min at 20 °C in four or five replicates. Results were recorded in terms of heat emitted (μ W) and recalculated per gram dry weight of sample, thus converted into specific heat production rate. The heat emitted by the bacteria themselves was not a significant factor in this study. Since 1.5 cm³ of the bacterial suspension emitted 60 μ W of heat, and since the amount of suspension used to inoculate each cotyledon was only 20 μ l, the amount of heat emitted by the bacteria present in each sample was negligible in comparison to the amount of heat emitted by the plant tissue itself.

2.3. Measurement of membrane permeability

After inoculation, the cotyledons were cut and placed in Petri dishes containing 10 cm³ of distilled water. Membrane permeability was determined by measuring ion leakage from the cotyledons with an OK-102/10 conductivity meter (Radelkis, Budapest, Hungary) in accordance with the method described by Barna et al. [10]. The first measurement was made 4 h after inoculation. Subsequent measurements were made 1, 2, 3, 6 and 7 days after infection.

3. Results

One day after inoculation, localized tissue necrosis was visible on the cotyledons. Two days after inoculation, the cotyledons were completely withered in the plants not pre-treated with BR₂₇. On the other hand, in infected plants pre-treated with BR₂₇, the spread of necrosis was significantly reduced and remained confined to a small area surrounding the injection site (Fig. 1). One day after inoculation, heat emission was significantly higher in infected cotyledons than in uninfected cotyledons (cultivar ‘Huzar’). This indicated that metabolic activity increased in tissue during the hypersensitive response induced by bacterial infection. In uninfected cotyledons, pre-treatment with BR₂₇ had no effect on heat emission. On the other hand, in infected cotyledons, pre-treatment with BR₂₇ significantly increased heat emission in both of the cultivars tested. This strongly suggests that BR₂₇ plays a role in preventing necrosis due to infection by incompatible bacteria in oilseed rape cotyledons (Fig. 2). Membrane permeability rapidly increased from the first to the second or third day after inoculation. In inoculated plants not pre-treated with BR₂₇, membrane permeability gradually increased throughout the 7 day observa-



Fig. 1. Effect of BR₂₇ on necrosis caused by infection with *Pseudomonas syringae* pv. *syringae* in the oilseed rape cultivar ‘Licosmos’ 7 days after inoculation. The healthy cotyledon on the left had been pre-treated with 100 nM BR₂₇, whereas the withered cotyledon on the right had not been pre-treated with BR₂₇.

tion period in both of the cultivars tested. In inoculated plants pre-treated with BR₂₇, membrane permeability was significantly reduced starting on the second or third day after inoculation and continuing for several days in both of the cultivars tested, espe-




Fig. 2. Specific heat production rate in cotyledons of the oilseed rape cultivars Huzar (A) and Licosmos (B) 1 day after inoculation with *Pseudomonas syringae* pv. *syringae* both with and without pre-treatment with 100 nM BR₂₇. Abbreviations: control, cotyledons pre-treated with water; BR₂₇, cotyledons pre-treated with BR₂₇ (100 nM) solution; Ps.s., cotyledons injected with suspension of incompatible bacteria *P. syringae* pv. *syringae* (10^8 cfu cm⁻³); BR₂₇ Ps.s., cotyledons pre-treated with BR₂₇ and injected with *P. syringae* pv. *syringae*; DW, dry weight. Data marked with the same letter are not significantly different according to Duncan's multiple-range *t*-test at *P* < 0.05.

Download English Version:

<https://daneshyari.com/en/article/675648>

Download Persian Version:

<https://daneshyari.com/article/675648>

[Daneshyari.com](https://daneshyari.com)