


Accepted Manuscript

Is there a general relationship between the exergy and HHV for rice residues?

Yaning Zhang, Qian Wang, Bingxi Li, Hongtao Li, Wenke Zhao



PII: S0960-1481(17)30984-9
DOI: 10.1016/j.renene.2017.10.022
Reference: RENE 9310
To appear in: *Renewable Energy*
Received Date: 20 February 2017
Revised Date: 03 September 2017
Accepted Date: 07 October 2017

Please cite this article as: Yaning Zhang, Qian Wang, Bingxi Li, Hongtao Li, Wenke Zhao, Is there a general relationship between the exergy and HHV for rice residues?, *Renewable Energy* (2017), doi: 10.1016/j.renene.2017.10.022

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

A specific relationship between exergy and HHV is observed for rice husks.

A specific relationship between exergy and HHV is observed for rice straws.

A general relationship between exergy and HHV is observed for rice residues.

The general relationship can be used for expeditious estimation of the exergy of rice residues.

Download English Version:

<https://daneshyari.com/en/article/6764952>

Download Persian Version:

<https://daneshyari.com/article/6764952>

[Daneshyari.com](https://daneshyari.com)