


Accepted Manuscript

Title: Self-harm and suicide attempts in Schizophrenia

Authors: Kiran Jakhar, Ram Pratap Beniwal, Triptish Bhatia,
Smita N. Deshpande

PII: S1876-2018(17)30083-7
DOI: <http://dx.doi.org/10.1016/j.ajp.2017.08.012>
Reference: AJP 1221



To appear in:

Received date: 7-2-2017
Revised date: 9-8-2017
Accepted date: 13-8-2017

Please cite this article as: Jakhar, Kiran, Beniwal, Ram Pratap, Bhatia, Triptish, Deshpande, Smita N., Self-harm and suicide attempts in Schizophrenia. Asian Journal of Psychiatry <http://dx.doi.org/10.1016/j.ajp.2017.08.012>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title Page

Self-harm and suicide attempts in Schizophrenia

Author list: Kiran Jakhar¹, Ram Pratap Beniwal¹, Triptish Bhatia², Smita N. Deshpande¹

¹Dept. Of Psychiatry, Centre of Excellence in Mental Health, PGIMER-Dr. R.M.L. Hospital, New Delhi, kjakhar48@gmail.com; beniwal_9@yahoo.co.in;

²GRIP- NIH Project, Dept. of Psychiatry, Centre of Excellence in Mental Health, PGIMER-Dr. R.M.L. Hospital, New Delhi, bhatiatriptish@yahoo.co.in

Corresponding Author:

Prof. Smita N. Deshpande

Department of Psychiatry,

Centre of Excellence in Mental Health

PGIMER- Dr. Ram Manohar Lohia Hospital, New Delhi

Email: smitadeshp@gmail.com

HIGHLIGHTS

- One fourth of schizophrenia patients reported self-harm including suicide attempt.
- Prevalence of suicide attempts was 10% in the sample.
- Fifty percent of attempted suicide was within one year of onset of illness
- Current age and past month GAF were significantly correlated with suicide attempt.
- Insecticide or overdose of medication was most common method of suicide attempt.

Self-harm and suicide attempts in Schizophrenia

Abstract

Background: The risk of suicide among persons with schizophrenia (SZ) is higher than in general population, with multiple contributory factors. We assessed the prevalence of risk of deliberate self-harm and suicide attempts, along with associated socio-demographic and clinical

Download English Version:

<https://daneshyari.com/en/article/6787714>

Download Persian Version:

<https://daneshyari.com/article/6787714>

[Daneshyari.com](https://daneshyari.com)