

Mice that are resistant to diet-induced weight loss have greater food anticipatory activity and altered melanocortin-3 receptor (MC3R) and Dopamine receptor 2 (D2) gene expression

L.M. Vaanholt, Sharon E. Mitchell, Rachel E. Sinclair, J.R. Speakman

PII: S0018-506X(15)00114-2
DOI: doi: 10.1016/j.yhbeh.2015.06.006
Reference: YHBEH 3897

To appear in: *Hormones and Behavior*

Received date: 18 March 2015
Revised date: 12 June 2015
Accepted date: 13 June 2015

Please cite this article as: Vaanholt, L.M., Mitchell, Sharon E., Sinclair, Rachel E., Speakman, J.R., Mice that are resistant to diet-induced weight loss have greater food anticipatory activity and altered melanocortin-3 receptor (MC3R) and Dopamine receptor 2 (D2) gene expression, *Hormones and Behavior* (2015), doi: 10.1016/j.yhbeh.2015.06.006

Mice that are resistant to diet-induced weight loss have greater food anticipatory activity and altered melanocortin-3 receptor (MC3R) and Dopamine receptor 2 (D2) gene expression.

L.M. Vaanholt[1], Sharon E. Mitchell[1], Rachel E. Sinclair[1], and J.R. Speakman[1,2]

1. Institute of Biological and Environmental Sciences, University of Aberdeen, Aberdeen, UK
2. State key laboratory of Molecular Developmental Biology, Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing, China

Corresponding author:

Dr. Lobke Maria Vaanholt
Institute of Biological and Environmental Sciences
University of Aberdeen
Zoology Building
Aberdeen AB24 2TZ
United Kingdom
Tel: +44 (0)1224 27 3255
Fax: +44 (0)1224 27 2396
E-mail: l.vaanholt@abdn.ac.uk

ABSTRACT

Diet-induced weight loss varies considerable between individuals, but the mechanisms driving these individual differences remain largely unknown. Here we