


Accepted Manuscript



Early Life Stress and Trauma and Enhanced Limbic Activation to Emotionally Valenced Faces in Depressed and Healthy Children

Hideo Suzuki , PhD Joan L. Luby , MD Kelly N. Botteron , MD Rachel Dietrich , BA
Mark P. McAvoy , PhD Deanna M. Barch , PhD

PII: S0890-8567(14)00295-0

DOI: [10.1016/j.jaac.2014.04.013](https://doi.org/10.1016/j.jaac.2014.04.013)

Reference: JAAC 1000

To appear in: *Journal of the American Academy of Child & Adolescent Psychiatry*

Received Date: 22 July 2013

Revised Date: 1 April 2014

Accepted Date: 3 April 2014

Please cite this article as: Suzuki H, Luby JL, Botteron KN, Dietrich R, McAvoy MP, Barch DM, Early Life Stress and Trauma and Enhanced Limbic Activation to Emotionally Valenced Faces in Depressed and Healthy Children, *Journal of the American Academy of Child & Adolescent Psychiatry* (2014), doi: 10.1016/j.jaac.2014.04.013.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Early Life Stress and Trauma and Enhanced Limbic Activation to Emotionally Valenced Faces in Depressed and Healthy Children

Running head: Stress and Functional Activation

Hideo Suzuki, PhD, Joan L. Luby, MD, Kelly N. Botteron, MD, Rachel Dietrich, BA, Mark P. McAvoy, PhD, and Deanna M. Barch, PhD

Clinical guidance is available at the end of this article.

Supplemental material cited in this article is available online.

Accepted May 1, 2014

This article was reviewed under and accepted by deputy editor Ellen Leibenluft, MD.

Drs. Suzuki, Luby, Botteron, McAvoy, Barch, and Ms. Dietrich are with Washington University in St. Louis, MO.

This study was supported by National Institutes of Health (NIH) grants MH064769 (J.L.L.) and MH090786 (J.L.L., D.M.B., K.N.B.).

The authors would like to thank the families and children who participated in these studies, and special thanks to staff members at the Early Emotional Development Program (EEDP) at Washington University in St. Louis for their assistance.

Disclosure: Dr. Luby has received grant or research support from the National Institute of Mental Health (NIMH), the Communities Healing Adolescent Depression and Suicide (CHADS) Coalition, and the Sydney R. Baer Foundation. Dr. Botteron has received grant or research support from the Eunice Kennedy Shriver National Institute of Child Health and Human Development, the National Institute of Biomedical Imaging and Bioengineering, NIMH, and Autism Speaks. Dr. Barch has received grant or research support from NIMH, the Brain & Behavior Research Foundation, and NIH. She has also served as a consultant for Pfizer, Roche, and Angen on schizophrenia-related work. Drs. Suzuki, McAvoy, and Ms. Dietrich report no biomedical financial interests or potential conflicts of interest.

Correspondence to: Hideo Suzuki, Washington University School of Medicine, Department of Psychiatry, Campus Box 8134, 660 Euclid Ave., Saint Louis, MO 63110, USA; email: suzukih@psychiatry.wustl.edu, hsuzuki@laureateinstitute.org

Download English Version:

<https://daneshyari.com/en/article/6797133>

Download Persian Version:

<https://daneshyari.com/article/6797133>

[Daneshyari.com](https://daneshyari.com)