

## Accepted Manuscript

Use of antipsychotic blood levels in clinician decision making: A cross-over study using clinical vignettes of patients with schizophrenia

Adam Savitz , Rama Melkote , Ralph Riley , Maria A Pobre , Kelly McQuarrie , David Williamson , Benjamin Banderas

PII: S0165-1781(17)31680-3  
DOI: [10.1016/j.psychres.2018.05.014](https://doi.org/10.1016/j.psychres.2018.05.014)  
Reference: PSY 11403



To appear in: *Psychiatry Research*

Received date: 13 September 2017  
Revised date: 20 March 2018  
Accepted date: 1 May 2018

Please cite this article as: Adam Savitz , Rama Melkote , Ralph Riley , Maria A Pobre , Kelly McQuarrie , David Williamson , Benjamin Banderas , Use of antipsychotic blood levels in clinician decision making: A cross-over study using clinical vignettes of patients with schizophrenia, *Psychiatry Research* (2018), doi: [10.1016/j.psychres.2018.05.014](https://doi.org/10.1016/j.psychres.2018.05.014)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Highlights**

- CASP was developed to capture clinical decision making in patients with schizophrenia
- Providing ABLs under descriptive conditions were used to evaluate treatment decisions
- Clinicians responded they were more confident in their decision making using ABLs

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/6811150>

Download Persian Version:

<https://daneshyari.com/article/6811150>

[Daneshyari.com](https://daneshyari.com)