

Accepted Manuscript

Cholecystokinin (CCK) level is higher among first time suicide attempters than healthy controls, but is not associated with higher depression scores

Leila Jahangard , Rahmat Solgy , Iraj Salehi ,
Seyed Kazem Taheri , Edith Holsboer-Trachsler ,
Mohammad Haghighi , Serge Brand

PII: S0165-1781(18)30167-7
DOI: [10.1016/j.psychres.2018.05.031](https://doi.org/10.1016/j.psychres.2018.05.031)
Reference: PSY 11420



To appear in: *Psychiatry Research*

Received date: 25 January 2018
Revised date: 26 April 2018
Accepted date: 11 May 2018

Please cite this article as: Leila Jahangard , Rahmat Solgy , Iraj Salehi , Seyed Kazem Taheri , Edith Holsboer-Trachsler , Mohammad Haghighi , Serge Brand , Cholecystokinin (CCK) level is higher among first time suicide attempters than healthy controls, but is not associated with higher depression scores, *Psychiatry Research* (2018), doi: [10.1016/j.psychres.2018.05.031](https://doi.org/10.1016/j.psychres.2018.05.031)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

HIGHLIGHTS

- We assessed suicide attempters within the first 12h after suicide attempt
- Cholecystokinin (CCK) levels and depression scores were higher among suicide attempters
- Statistically, CCK levels and depression scores were unrelated

ACCEPTED MANUSCRIPT

Download English Version:

<https://daneshyari.com/en/article/6811200>

Download Persian Version:

<https://daneshyari.com/article/6811200>

[Daneshyari.com](https://daneshyari.com)