

Accepted Manuscript

More intensive probing increases the detection of social anxiety disorders in schizophrenia

Marc-André Roy MD, FRCP , Chantal Vallières B.Psy ,
Catherine Lehoux PhD , Louis-David Leclerc MD ,
Marie-France Demers B Pharm, MSC, BCPP ,
Amélie M. Achim PhD

PII: S0165-1781(18)30741-8
DOI: [10.1016/j.psychres.2018.07.032](https://doi.org/10.1016/j.psychres.2018.07.032)
Reference: PSY 11581



To appear in: *Psychiatry Research*

Received date: 24 April 2018
Revised date: 12 July 2018
Accepted date: 18 July 2018

Please cite this article as: Marc-André Roy MD, FRCP , Chantal Vallières B.Psy , Catherine Lehoux PhD , Louis-David Leclerc MD , Marie-France Demers B Pharm, MSC, BCPP , Amélie M. Achim PhD , More intensive probing increases the detection of social anxiety disorders in schizophrenia, *Psychiatry Research* (2018), doi: [10.1016/j.psychres.2018.07.032](https://doi.org/10.1016/j.psychres.2018.07.032)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- It is important to identify the source of social anxiety in psychotic patients;
- Treating co-morbid social anxiety disorders (SAD) may support recovery in psychosis;
- Variations in the diagnostic methods may affect the detection of SAD;
- More intensive probing for SAD increases its prevalence estimate;
- Psychotic patients with SAD may have distinct clinical profiles.

Download English Version:

<https://daneshyari.com/en/article/6811300>

Download Persian Version:

<https://daneshyari.com/article/6811300>

[Daneshyari.com](https://daneshyari.com)