## **Accepted Manuscript**

Nightmares and Suicide Risk in Psychiatric Patients: The Roles of Hopelessness and Male Depressive Symptoms

Dorian A. Lamis, Marco Innamorati, Denise Erbuto, Isabella Berardelli, Franco Montebovi, Gianluca Serafini, Mario Amore, Barry Krakow, Paolo Girardi, Maurizio Pompili

PII: S0165-1781(17)32005-X

DOI: 10.1016/j.psychres.2018.03.053

Reference: PSY 11285

To appear in: Psychiatry Research

Received date: 29 October 2017 Revised date: 13 March 2018 Accepted date: 22 March 2018



Please cite this article as: Dorian A. Lamis, Marco Innamorati, Denise Erbuto, Isabella Berardelli, Franco Montebovi, Gianluca Serafini, Mario Amore, Barry Krakow, Paolo Girardi, Maurizio Pompili, Nightmares and Suicide Risk in Psychiatric Patients: The Roles of Hopelessness and Male Depressive Symptoms, *Psychiatry Research* (2018), doi: 10.1016/j.psychres.2018.03.053

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Highlight

- Nightmares play an important role in well-being and may be important for outcome of psychiatric disorders
- Hopelessness and depression have a contributing role in the increasing suicide risk
- We found that patients with more nightmares had more hopelessness and depression as well as suicide risk
- We conclude that the assessment of nightmares help clinicians in defining the complexity of each patients and provide the best treatment



## Download English Version:

## https://daneshyari.com/en/article/6811432

Download Persian Version:

https://daneshyari.com/article/6811432

<u>Daneshyari.com</u>