

## Accepted Manuscript

Facial emotion recognition, socio-occupational functioning and expressed emotions in schizophrenia versus bipolar disorder"

Umesh Thonse , Rishikesh V. Behere , Samir Kumar Praharaj ,  
Podila Sathya Venkata Narasimha Sharma

PII: S0165-1781(17)31427-0  
DOI: [10.1016/j.psychres.2018.03.027](https://doi.org/10.1016/j.psychres.2018.03.027)  
Reference: PSY 11259



To appear in: *Psychiatry Research*

Received date: 1 August 2017  
Revised date: 5 January 2018  
Accepted date: 9 March 2018

Please cite this article as: Umesh Thonse , Rishikesh V. Behere , Samir Kumar Praharaj , Podila Sathya Venkata Narasimha Sharma , Facial emotion recognition, socio-occupational functioning and expressed emotions in schizophrenia versus bipolar disorder", *Psychiatry Research* (2018), doi: [10.1016/j.psychres.2018.03.027](https://doi.org/10.1016/j.psychres.2018.03.027)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Highlights**

- Ours is one of the first studies to examine the association between facial emotion recognition and expressed emotions between two groups of severe mental illness in the remission phase.
- Used Indian made facial emotion recognition test called TRENDS (Tools for Recognition of Emotions in Neuropsychiatric Disorders)
- Found no difference in emotion recognition abilities between patients with schizophrenia and bipolar disorder.
- Eventhough, patients with schizophrenia were in remission phase of the illness they were experiencing more critical comments from their caregivers than patients with bipolar disorder.
- Patients with schizophrenia in current study were having poorer socio occupational functioning with comparison to patients with bipolar disorder in the current study.
- We have noticed significant positive correlation between TRACS score and socio occupational functioning impairment in patients with bipolar disorder.
- Interestingly this present study has found that impairment in emotion recognition in patients with bipolar disorder associated with hostility and dissatisfaction from their primary caregiver.

Download English Version:

<https://daneshyari.com/en/article/6811664>

Download Persian Version:

<https://daneshyari.com/article/6811664>

[Daneshyari.com](https://daneshyari.com)