

Psychotic symptoms and quality of life: a mediation analysis of daily-life coping

Emilio López-Navarro, Cristina del Canto, Antoni Mayol, Ovidio Fernández-Alonso, Enric Munar



PII: S0165-1781(17)30819-3

DOI: <http://dx.doi.org/10.1016/j.psychres.2017.09.034>

Reference: PSY10845

To appear in: *Psychiatry Research*

Received date: 12 May 2017

Accepted date: 12 September 2017

Cite this article as: Emilio López-Navarro, Cristina del Canto, Antoni Mayol, Ovidio Fernández-Alonso and Enric Munar, Psychotic symptoms and quality of life: a mediation analysis of daily-life coping, *Psychiatry Research*, <http://dx.doi.org/10.1016/j.psychres.2017.09.034>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Psychotic symptoms and quality of life: a mediation analysis of daily-life coping**

Emilio López-Navarro<sup>a1</sup>, Cristina del Canto<sup>b</sup>, Antoni Mayol<sup>a,c</sup>, Ovidio Fernández-Alonso<sup>c</sup>, Enric Munar<sup>a</sup>

<sup>a</sup>EvoCog Group – Associated Unit to CSIC, University of Balearic Islands, Balearic Islands, Spain.

<sup>b</sup>Department of Clinical Psychology – Son Espases Hospital. Balearic Health Service. Balearic Islands. Spain

<sup>c</sup>UCR Serralta Community Rehabilitation Centre. Balearic Health Service. Balearic Islands. Spain

Emilio.lopez@uib.es

### Abstract

Prior studies suggest the relationship between psychotic symptoms and Quality of Life (QoL) may be mediated by diverse constructs. QoL in schizophrenia-related disorders has been related with coping with daily stressors. Based on previous studies, our hypothesis was that coping mediates the relationship between psychotic symptoms and QoL. Therefore, the aim of the study was to test the hypothesis in a sample of people with schizophrenia-related disorders from a community rehabilitation center. Sixty-six patients were assessed using PANSS, WHOQOL-BREF questionnaire, and COPE Inventory. Regression

---

<sup>1</sup> EvoCog Group, IFISC, Associated Unit to CSIC, Guillem Cifre Building, University of Balearic Islands. Ctra Valldemossa km 7,5. Balearic Islands, Spain. Phone: +34 971 259888

Download English Version:

<https://daneshyari.com/en/article/6811792>

Download Persian Version:

<https://daneshyari.com/article/6811792>

[Daneshyari.com](https://daneshyari.com)