

# Author's Accepted Manuscript

The association between rehabilitation programs and metabolic syndrome in chronic inpatients with schizophrenia

Yi-Chun Lin, Chien-Liang Lai, Hung-Yu Chan



PII: S0165-1781(17)30485-7  
DOI: <https://doi.org/10.1016/j.psychres.2017.11.081>  
Reference: PSY11028

To appear in: *Psychiatry Research*

Received date: 18 March 2017  
Revised date: 4 October 2017  
Accepted date: 26 November 2017

Cite this article as: Yi-Chun Lin, Chien-Liang Lai and Hung-Yu Chan, The association between rehabilitation programs and metabolic syndrome in chronic inpatients with schizophrenia, *Psychiatry Research*, <https://doi.org/10.1016/j.psychres.2017.11.081>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**The association between rehabilitation programs and metabolic syndrome in chronic inpatients with schizophrenia**

Yi-Chun Lin<sup>a</sup>, Chien-Liang Lai<sup>a</sup>, Hung-Yu Chan, M.D.<sup>a,b\*</sup>

<sup>a</sup>Department of General Psychiatry, Taoyuan Psychiatric Center, Ministry of Health and Welfare, Taoyuan, Taiwan

<sup>b</sup>Department of Psychiatry, National Taiwan University Hospital and College of Medicine, National Taiwan University, Taipei, Taiwan

Yi-Chun Lin and Chien-Liang Lai contributed equally as the first author of this article.

\*Correspondence and reprint requests to: Hung-Yu Chan, M.D., PhD, , Taoyuan Psychiatric Center, Ministry of Health and Welfare, No. 71, Long-show Street, Taoyuan City 33058, Taiwan. e-mail: jan30@seed.net.tw. tel.: +886-(3)-369-8553; fax: +886-(3)-369-9498.

**Abstract**

The correlation between different rehabilitation programs and the prevalence of metabolic syndrome in people with schizophrenia is unclear. We tested the association in chronic inpatients with schizophrenia of a psychiatric hospital in Taiwan. Patients with schizophrenia and age from 20 to 65 years old were included. The criteria of metabolic syndrome were according to the adapted Adult Treatment Protocol for

Download English Version:

<https://daneshyari.com/en/article/6811901>

Download Persian Version:

<https://daneshyari.com/article/6811901>

[Daneshyari.com](https://daneshyari.com)