

## Author's Accepted Manuscript

Efficacy of bright light therapy in bipolar depression

Nese Yorguner Kupeli, Serkut Bulut, Gresa Carkaxhiu Bulut, Emel Kurt, Kaan Kora



PII: S0165-1781(17)31113-7  
DOI: <https://doi.org/10.1016/j.psychres.2017.12.020>  
Reference: PSY11056

To appear in: *Psychiatry Research*

Received date: 17 June 2017  
Revised date: 23 October 2017  
Accepted date: 9 December 2017

Cite this article as: Nese Yorguner Kupeli, Serkut Bulut, Gresa Carkaxhiu Bulut, Emel Kurt and Kaan Kora, Efficacy of bright light therapy in bipolar depression, *Psychiatry Research*, <https://doi.org/10.1016/j.psychres.2017.12.020>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# Efficacy of bright light therapy in bipolar depression

Nese Yorguner Kupeli<sup>\*1</sup>, Serkut Bulut<sup>2</sup>, Gresa Carkaxhiu Bulut<sup>3</sup>, Emel Kurt<sup>4</sup>, Kaan Kora<sup>5</sup>

<sup>1</sup>Department of Psychiatry, Marmara University Pendik Training and Research Hospital, Istanbul, Turkey,

<sup>2</sup>Department of Psychiatry, Mus State Hospital, Mus, Turkey,

<sup>3</sup>Department of Child and Adolescent Psychiatry, Mus State Hospital, Mus, Turkey,

<sup>4</sup>Department of Psychiatry, Hisar Intercontinental Hospital, Istanbul, Turkey,

<sup>5</sup>Private Practice, Vizyon Psychiatry, Istanbul, Turkey,

[nesevorguner@gmail.com](mailto:nesevorguner@gmail.com)

[serkutbulut@gmail.com](mailto:serkutbulut@gmail.com)

[gresacarkaxhiu@gmail.com](mailto:gresacarkaxhiu@gmail.com)

[emelbuyrazkurt@gmail.com](mailto:emelbuyrazkurt@gmail.com)

[kaankora@gmail.com](mailto:kaankora@gmail.com)

\***Correspondence:** Nese Yorguner Kupeli , **Address:** Department of Psychiatry, Marmara University Pendik Training and Research Hospital, Ustkaynarca, Pendik, Istanbul, Turkey. **Fax:** +90 0216 625 46 39

## ABSTRACT

For 30 years, bright light therapy (BLT) has been considered as an effective, well-tolerated treatment for seasonal affective disorder (SAD). Because of low response rates, new treatment strategies are needed for bipolar depression (BD), which resembles SAD in certain respects. Few placebo-controlled studies of BLT efficacy have been carried out for BD. Accordingly, this study evaluates the efficacy and safety of BLT as an add-on treatment for BD. Thirty-two BD outpatients were randomly assigned to BLT (10000 lux) or dim light (DL, <500 lux). During a two-week period, light was administered each morning for 30 min. The Hamilton Rating Scale for Depression and the Montgomery-Åsberg Depression Rating Scale assessed clinical outcome, and the UKU Side Effects Rating Scale evaluated side effects. No significant difference was observed in baseline

Download English Version:

<https://daneshyari.com/en/article/6812024>

Download Persian Version:

<https://daneshyari.com/article/6812024>

[Daneshyari.com](https://daneshyari.com)