Author's Accepted Manuscript

Factors Related to Physical Violence Experienced by Parents of Persons with Schizophrenia in Japan

Masako Kageyama, Phyllis Solomon, Sachiko Kita, Satoko Nagata, Keiko Yokoyama, Yukako Nakamura, Sayaka Kobayashi, Chiyo Fujii



www.elsevier.com/locate/psvchres

PII: S0165-1781(15)30595-3

DOI: http://dx.doi.org/10.1016/j.psychres.2016.06.036

PSY9787 Reference:

To appear in: Psychiatry Research

Received date: 30 October 2015 Revised date: 21 June 2016 Accepted date: 23 June 2016

Cite this article as: Masako Kageyama, Phyllis Solomon, Sachiko Kita, Satoko Nagata, Keiko Yokoyama, Yukako Nakamura, Sayaka Kobayashi and Chiyo Fujii, Factors Related to Physical Violence Experienced by Parents of Person Schizophrenia Japan, Psychiatry with in Research http://dx.doi.org/10.1016/j.psychres.2016.06.036

This is a PDF file of an unedited manuscript that has been accepted fo publication. As a service to our customers we are providing this early version o the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain

ACCEPTED MANUSCRIPT

Factors Related to Physical Violence Experienced by Parents of Persons with

Schizophrenia in Japan

Masako Kageyama^{a*}, Phyllis Solomon^{b1}, Sachiko Kita^a, Satoko Nagata^a, Keiko Yokoyama^{c2}, Yukako Nakamura^{d3}, Sayaka Kobayashi^{e4}, Chiyo Fujii^{e,4}

^aGraduate School of Medicine, the University of Tokyo

^bSchool of Social Policy & Practice, University of Pennsylvania, 3701 Locust Walk, Philadelphia, PA 19104-6214, USA

^cFaculty of Health Sciences, Saitama Prefectural University, 820 Sannomiya, Koshigaya City, Saitama 343-8540, JAPAN

^dNagoya University Graduate School of Medicine, 65 Tsurumai-Cho, Showa-ku, Nagoya, Aichi 466-8550, JAPAN

^eNational Institute of Mental Health, 4-1-1 Ogawahigashi-Cho, Kodaira City, Tokyo 187-8552, JAPAN

kageyama-tky@umin.ac.jp

solomonp@sp2.upenn.edu

sachiko-happygirl@hotmail.co.jp

satoko-tky@umin.ac.jp

yokoyama-keiko@spu.ac.jp

y-nakamu@med.nagoya-u.ac.jp

skobayashi@ncnp.go.jp

chyfujii@ncnp.go.jp

*Corresponding author: Department of Community Health Nursing, Graduate School of Medicine, the University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, JAPAN; Tel./fax: +81 3 5841 3649

Abstract

Most violence by patients with mental illness is perpetuated against family members rather

than the general public. However, there is insufficient research to reach a consensus on

¹ Tel: +1-215-898-5533, Fax: +1-215-573-2099

² Tel/Fax: +81-48-973-4769

³ Tel: +81-52-744-2282, Fax: +81-52-744-2293

⁴ Tel: +81-42-341-2711, Fax: +81-42-346-1944

Download English Version:

https://daneshyari.com/en/article/6812912

Download Persian Version:

https://daneshyari.com/article/6812912

<u>Daneshyari.com</u>