


Author's Accepted Manuscript

Psychotic-like experiences and disordered eating in the English general population

Ai Koyanagi, Andrew Stickley, Josep Maria Haro



PII: S0165-1781(15)30347-4
DOI: <http://dx.doi.org/10.1016/j.psychres.2016.04.045>
Reference: PSY9608

To appear in: *Psychiatry Research*

Received date: 15 September 2015
Revised date: 10 March 2016
Accepted date: 15 April 2016

Cite this article as: Ai Koyanagi, Andrew Stickley and Josep Maria Haro
Psychotic-like experiences and disordered eating in the English general population, *Psychiatry Research*
<http://dx.doi.org/10.1016/j.psychres.2016.04.045>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Psychotic-like experiences and disordered eating in the English general population

Ai Koyanagi^{a,b,*}, Andrew Stickley^c, Josep Maria Haro^{a,b}

^aParc Sanitari Sant Joan de Déu, Universitat de Barcelona, Fundació Sant Joan de Déu, Dr Antoni Pujades, 42, Sant Boi de Llobregat, Barcelona, 08830, Spain

^bInstituto de Salud Carlos III, Centro de Investigación Biomédica en Red de Salud Mental, CIBERSAM, Monforte de Lemos 3-5 Pabellón 11, 28029, Madrid

^cThe Stockholm Centre for Health and Social Change (SCOHOST), Södertörn University, Huddinge 141 89, Sweden

***Corresponding author.** Ai Koyanagi MD MSc PhD. Research and Development Unit, Parc Sanitari Sant Joan de Déu, Fundació Sant Joan de Déu, Dr. Antoni Pujades, 42, Sant Boi de Llobregat, Barcelona, 08830, Spain. Tel: (+34)936002685. FAX: (+34)936305319. E-mail: a.koyanagi@pssjd.org

Abstract

There are no studies on psychotic-like experiences (PLEs) and disordered eating in the general population. We aimed to assess this association in the English adult population. Data from the 2007 Adult Psychiatric Morbidity Survey (APMS) were analyzed. This was a nationally representative survey comprising 7403 English adults aged ≥ 16 years. The Psychosis Screening Questionnaire was used to identify the past 12-month occurrence of five forms of psychotic symptoms. Questions from the five-item SCOFF screening instrument were used to identify those with eating disorder (ED) symptoms and possible ED in the past year. The prevalence of any PLE was 5.1% (female) and 5.4% (male), while that of possible ED was 9.0% (female) and 3.5% (male). After adjustment for potential confounders, possible

Download English Version:

<https://daneshyari.com/en/article/6813010>

Download Persian Version:

<https://daneshyari.com/article/6813010>

[Daneshyari.com](https://daneshyari.com)