

## Author's Accepted Manuscript

Serum soluble urokinase-type plasminogen activator receptor levels in male patients with acute exacerbation of schizophrenia

Abdullah Genc, Tevfik Kalelioğlu, Nesrin Karamustafalioglu, Akif Tasdemir, Esra Sena Genc, Mustafa Akkus, Murat Emul



PII: S0165-1781(15)30106-2  
DOI: <http://dx.doi.org/10.1016/j.psychres.2016.01.018>  
Reference: PSY9378

To appear in: *Psychiatry Research*

Received date: 5 August 2015  
Revised date: 31 December 2015  
Accepted date: 8 January 2016

Cite this article as: Abdullah Genc, Tevfik Kalelioğlu, Nesrin Karamustafalioglu Akif Tasdemir, Esra Sena Genc, Mustafa Akkus and Murat Emul, Serum soluble urokinase-type plasminogen activator receptor levels in male patients with acute exacerbation of schizophrenia, *Psychiatry Research* <http://dx.doi.org/10.1016/j.psychres.2016.01.018>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain

Serum soluble urokinase-type plasminogen activator receptor levels in male patients with acute exacerbation of schizophrenia

Abdullah Genc <sup>a,\*</sup>, Tevfik Kaleliođlu <sup>b</sup>, Nesrin Karamustafalioglu <sup>b</sup>, Akif Tasdemir <sup>b</sup>, Esra Sena Genc <sup>b</sup>, Mustafa Akkus <sup>b</sup>, Murat Emul <sup>c</sup>

MDs, Departments of Psychiatry, <sup>a</sup> ŐiŐli Etfal Training and Education Hospital, <sup>b</sup> Bakırkoy Research and Training Hospital for Neurology and Mental Health, <sup>c</sup> Medical School of Cerrahpasa, İstanbul, TURKEY

\*Corresponding author: Abdullah Genc, MD,  
Department of Psychiatry,  
ŐiŐli Etfal Training and Education Hospital, Halaskargazi  
Avenue, Etfal Street, 34371, Istanbul, Turkey  
Tel: +90 212 543 65 65  
e-mail: [abdullahgenc@mail.com](mailto:abdullahgenc@mail.com)

## Abstract

Inflammatory abnormalities have been shown in the pathogenesis of schizophrenia. Soluble urokinase-type plasminogen activator receptor (suPAR) is a protein that is measurable in the circulating blood and reflects the inflammation in the body. We aimed to investigate serum suPAR levels in patients with schizophrenia who were in acute state and to compare with healthy controls. Forty five patients and 43 healthy controls were included in the study. We found no significant difference in suPAR levels between patients and controls, suggesting

Download English Version:

<https://daneshyari.com/en/article/6813663>

Download Persian Version:

<https://daneshyari.com/article/6813663>

[Daneshyari.com](https://daneshyari.com)