

## Accepted Manuscript

Title: Glucocorticoid receptor gene methylation moderates the association of childhood trauma and cortisol stress reactivity

Authors: Nina Alexander, Clemens Kirschbaum, Matthis Wankerl, Benjamin J. Stauch, Tobias Stalder, Susann Steudte-Schmiedgen, Markus Muehlhan, Robert Miller



PII: S0306-4530(17)31355-0  
DOI: <https://doi.org/10.1016/j.psyneuen.2018.01.020>  
Reference: PNEC 3825

To appear in:

Received date: 12-9-2017  
Revised date: 18-1-2018  
Accepted date: 22-1-2018

Please cite this article as: Alexander, Nina, Kirschbaum, Clemens, Wankerl, Matthis, Stauch, Benjamin J., Stalder, Tobias, Steudte-Schmiedgen, Susann, Muehlhan, Markus, Miller, Robert, Glucocorticoid receptor gene methylation moderates the association of childhood trauma and cortisol stress reactivity. *Psychoneuroendocrinology* <https://doi.org/10.1016/j.psyneuen.2018.01.020>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# Glucocorticoid receptor gene methylation moderates the association of childhood trauma and cortisol stress reactivity

**Abbreviated Title:** Glucocorticoid receptor methylation, trauma and cortisol reactivity

Nina Alexander<sup>1</sup>, Clemens Kirschbaum<sup>2</sup>, Matthis Wankerl<sup>2</sup>, Benjamin J. Stauch<sup>2</sup>, Tobias Stalder<sup>3</sup>, Susann Steudte-Schmiedgen<sup>2</sup>, Markus Muehlhan<sup>1</sup>, & Robert Miller<sup>2</sup>

<sup>1</sup> Department of Psychology, Faculty of Human Sciences, Medical School Hamburg, Hamburg, Germany,

<sup>2</sup> Department of Psychology, Faculty of Science, Technische Universität Dresden, Germany

<sup>3</sup> Clinical Psychology, University of Siegen, Siegen, Germany

## Word Count

Abstract: 284

Manuscript: 4829

Tables: 2

Figures: 2

Supplemental Material: 2

## Corresponding Author

Prof. Dr. Nina Alexander  
MSH Medical School Hamburg  
Department of Psychology, Faculty of Human Sciences  
Am Kaiserkai 1  
20457 Hamburg  
phone: +49 40 36122640  
e-mail: nina.alexander@medicalschooll-hamburg.de

## Highlights

- DNA methylation (DNA<sub>M</sub>) in the glucocorticoid receptor gene (*NR3C1-1F*) moderates the specific direction of HPA-axis dysregulation in childhood trauma survivors.
- Trauma survivors with increased *NR3C1-1F* DNA<sub>M</sub> displayed, on average, 10.4 nmol/l (62.3 %) higher peak cortisol levels during a laboratory stressor compared to those with low DNA<sub>M</sub>.
- In contrast, individuals who were unexposed or only moderately exposed to CT displayed a moderately sized cortisol stress response irrespective of *NR3C1-1F* methylation.
- Contrary to some studies work, our data provides no evidence for a direct association of childhood trauma and *NR3C1-1F* DNA<sub>M</sub> status.

## Abstract

Exposure to childhood trauma (CT) has been linked to sustained dysregulations of major stress response systems, including findings of both exaggerated and attenuated hypothalamus–pituitary–adrenal (HPA) axis activity. Likewise, CT constitutes a common risk factor for a broad range of

Download English Version:

<https://daneshyari.com/en/article/6817710>

Download Persian Version:

<https://daneshyari.com/article/6817710>

[Daneshyari.com](https://daneshyari.com)