



Short communication

Single dose testosterone administration reduces loss chasing in healthy females



Yin Wu^{a,b,1}, Jinting Liu^{a,b,c,1}, Lujing Qu^b, Christoph Eisenegger^d, Luke Clark^e, Xiaolin Zhou^{b,f,g,h,*}

^a Research Center for Brain Function and Psychological Science, Shenzhen University, Shenzhen, China

^b Center for Brain and Cognitive Sciences and Department of Psychology, Peking University, Beijing, China

^c China Center for Special Economic Zone Research, Shenzhen University, Shenzhen, China

^d Neuropsychopharmacology and Biopsychology Unit, Department of Basic Psychological Research and Research Methods, Faculty of Psychology, University of Vienna, Vienna, Austria

^e Centre for Gambling Research at UBC, Department of Psychology, University of British Columbia, Vancouver, British Columbia, Canada

^f Key Laboratory of Machine Perception (Ministry of Education), Peking University, Beijing, China

^g Beijing Key Laboratory of Behavior and Mental Health, Peking University, Beijing, China

^h IDG/McGovern Institute for Brain Research, Peking University, Beijing, China

ARTICLE INFO

Article history:

Received 22 February 2016

Received in revised form 3 May 2016

Accepted 3 May 2016

Keywords:

Testosterone

Gambling

Risk taking

Punishment sensitivity

ABSTRACT

Testosterone has been linked to modulation of impulsivity and risky choice, potentially mediated by changes in reward or punishment sensitivity. This study investigated the effect of testosterone on risk-taking and the adjustment of risk-taking on trials following a gain or a loss. Loss chasing is operationalized herein as the propensity to recover losses by increasing risky choice. Healthy female participants ($n = 26$) received a single-dose of 0.5 mg sublingual testosterone in a double-blind, placebo-controlled crossover design. At 240 min post-administration, participants performed a gambling task with a high and a low risk option. In the placebo condition, participants were more likely to choose the high risk option following losses compared to wins. This effect was abolished on the testosterone session. Ignoring prior outcomes, no overall changes in risk-taking were observed. Our data indicate that testosterone affects human decision-making via diminishing sensitivity to punishment.

© 2016 Published by Elsevier Ltd.

1. Introduction

Individuals tend to respond to losses during gambling by increasing their bets on subsequent gambles, a phenomenon referred as “loss chasing”. This tendency is considered a cardinal symptom of disordered gambling, and is one of the most commonly endorsed symptoms for gambling disorder in the Diagnostic and Statistical Manual of Mental Disorders (DSM) (Toce-Gerstein et al., 2003). In a large naturalistic dataset of online poker players, experienced players responded to major losses (>\$1,000 in a single hand) by becoming more aggressive on the next hand, and betting on weaker hands to stay in the game (Smith et al., 2009). Laboratory studies have shown increased risk taking behavior following a single loss compared with a single gain (Gehring and Willoughby,

2002). Loss chasing has also been demonstrated in a cumulative manner such that the probability of placing a high bet increased as losing streaks grew longer (Studer et al., 2015).

While there have been efforts to clarify the neurochemical basis of loss chasing with a particular focus on brain monoamines (Campbell-Meiklejohn et al., 2011), the role of hormones has not been systematically investigated. Evidence for a potential role of the sex hormone testosterone stems from research showing that males with higher levels of salivary testosterone showed higher levels of risk-taking according to self-reports (Vermeersch et al., 2008), on laboratory tasks with financial incentives (Apicella et al., 2008, 2014), and in naturalistic studies of stock traders (Coates and Herbert, 2008). Nevertheless, some inconsistent findings have also been reported, including evidence for nonlinearity (Stanton et al., 2011), effects of gender, and differences with incentive structure (see Apicella et al., 2015 for a review).

In studies manipulating testosterone levels, sublingual testosterone administration in women was shown to increase risky choices on the Iowa Gambling Task. Such an effect may reflect a

* Corresponding author at: Department of Psychology, Peking University, Beijing 100871, China.

E-mail address: xz104@pku.edu.cn (X. Zhou).

¹ These authors have contributed equally to this work.

reduction in punishment sensitivity, and/or an increase in reward sensitivity (van Honk et al., 2004). A trial of the aromatase inhibitor letrozole in healthy men showed that raised testosterone levels were associated with increased risk-taking under ambiguity (i.e., unknown probabilities) but not under conditions with known probabilities (Goudriaan et al., 2010).

The aim of the present study was to investigate the effects of a single dose of testosterone on loss chasing in women. We used a double-blind, placebo-controlled crossover design. Participants played a gambling task where they were asked to choose one of the two cards, displaying the numerals 5 and 25. Each card could win or lose the displayed amount with a probability of 0.5. Thus, the two options both have expected values of zero, but the 5 option is operationally “safe” (low outcome variance) and the 25 option is risky. We tested for an overall effect of testosterone administration to increase risk-taking on the task, and we hypothesized that ignoring prior outcome, testosterone would increase risk-taking. We also tested an effect on loss-chasing, manifested in how participants adjusted their choice towards the risky option following losses compared to wins (Gehring and Willoughby, 2002). We predicted that testosterone would influence loss-chasing given its effect upon reward/punishment sensitivity (van Honk et al., 2004). Second-digit-to-fourth digit (2D:4D) ratios were measured, in light of earlier findings that the effect of testosterone administration can be moderated by this proxy for prenatal testosterone exposure in both women (van Honk et al., 2011) and men (Carré et al., 2015).

2. Methods

2.1. Participants

Twenty-six healthy females (mean age = 21.5 years, $SD = 1.96$; age range = 18–25) were recruited through university advertisements. All the contacted persons were screened in a telephone interview to exclude individuals (5 in total) taking psychotropic medications, or having any psychiatric or neurological disorders. We only recruited females as the dosing and pharmacokinetics associated with single sublingual administration of testosterone are only established for women, and are unclear in men (Tuiten et al., 2000). Participants were instructed to abstain from alcohol, caffeine intake and smoking for 24 h before the testing session. They were tested within 10 days after the beginning of the menstrual cycle (interval of two testing days, $M = 25.96$ days, $SD = 6.32$, range = 11–36), when endogenous levels of sex hormones tend to be low and stable. Each participant received a single dose of testosterone and placebo in a crossover, double-blind, placebo-controlled, within-participant design. The study was conducted in accordance with the Declaration of Helsinki and was approved by the Ethics Committee of the Department of Psychology, Peking University. Written informed consent was obtained from all participants. Participants were paid 240 Chinese Yuan (~\$36.84) as a show-up fee. Then they were endowed with 20 Yuan to play the gambling task. The points that were gained or lost on the task were added or subtracted from this endowment, which was a bonus payment.

2.2. Testosterone administration

The testosterone preparation contained 0.5 mg of testosterone base, 5 mg of cyclodextrin (as a carrier), 5 mg of ethanol, and 5 mL of water. The placebo contained no testosterone but was otherwise identical. Testosterone and placebo were administered sublingually. All sessions started at 13:00 and lasted approximately 5 h. Due to the established time lag of 4 h for behavioral effects to appear after sublingual application of 0.5 mg testosterone in healthy young




Fig. 1. Sequence of events in a single trial. This trial displays a loss outcome, on which the participant has lost 25 on the chosen card (For interpretation of the references to color in the figure, the reader is referred to the web version of the article).

women (Tuiten et al., 2000), the gambling task commenced 4 h post-dosing. The participants also completed two further tasks of social cognition, not reported here. During the waiting period, participants rested in the laboratory and were provided with newspapers and magazines that were not related to the study.

2.3. Gambling task

Participants completed 480 experimental trials on a computerized gambling task modified from Gehring and Willoughby (2002); see Fig. 1), and programmed using Presentation software (Neurobehavioral System Inc.). On each trial, participants were presented with two cards, displaying the numerals 5 and 25 that corresponded to the possible gain or loss outcomes with a probability of 0.5. Participants were informed that each point corresponded to 0.1 Chinese Yuan they could earn or lose. But they were not informed of the gain or loss probability, although the outcomes were pseudo-randomized to ensure the effective expected value was zero. They were asked to choose one of them by pressing the corresponding button within 3000 ms. After a jittered blank screen, the outcome on the selected card was presented for 1000 ms. The + or – symbols on the card indicated the amounts the participant won or lost. The next trial started after a 800 ms intertrial interval.

2.4. Digit ratio measurement

Digit ratio was measured from an image scan of the right-hand by measuring the length of the index (2D) and ring (4D) finger from the ventral proximal crease to the tip of the finger using Adobe Photoshop. Two experienced raters (blind to the purpose of the study) measured the 2D:4D ratio on three occasions, and the mean value was used for analysis. Inter-rater reliability was high, $r = 0.96$, $p < 0.001$.

Download English Version:

<https://daneshyari.com/en/article/6818011>

Download Persian Version:

<https://daneshyari.com/article/6818011>

[Daneshyari.com](https://daneshyari.com)