


Author's Accepted Manuscript

Prevalence and risk factors of type 2 diabetes in major depression: A study on 703 individuals referred for sleep examinations

Matthieu Hein, Jean-Pol Lanquart, Gwénoél Loas, Philippe Hubain, Paul Linkowski



www.elsevier.com/locate/bios

PII: S0033-3182(17)30223-2

DOI: <https://doi.org/10.1016/j.psym.2017.11.003>[10.1016/j.psychres.2017.08.061](https://doi.org/10.1016/j.psychres.2017.08.061)

Reference: PSYM834

To appear *Psychosomatics*
in:

Cite this article as: Matthieu Hein, Jean-Pol Lanquart, Gwénoél Loas, Philippe Hubain and Paul Linkowski, Prevalence and risk factors of type 2 diabetes in major depression: A study on 703 individuals referred for sleep examinations, *Psychosomatics*, doi:10.1016/j.psym.2017.11.003

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Prevalence and risk factors of type 2 diabetes in major depression: A study on 703 individuals referred for sleep examinations.

Matthieu Hein^a, Jean-Pol Lanquart^a, Gwéno^lé Loas^a, Philippe Hubain^a and Paul Linkowski^a

^aErasme Hospital, Department of Psychiatry and Sleep Laboratory, Université libre de Bruxelles, ULB, Brussels, Belgium.

Route de Lennik, 808 - 1070 Anderlecht - Belgium

Phone : 0032/25553741 – FAX : 0032/25556955

E-mail : SecMed.Psy@erasme.ulb.ac.be – matthieu.hein@erasme.ulb.ac.be

Abstract

Background: Several studies have investigated the prevalence and risk factors of depression in individuals with type 2 diabetes. However, few studies have investigated the prevalence and risk factors for type 2 diabetes in major depression. The aim of this study was to examine the prevalence and risk factors of type 2 diabetes in a large sample of individuals with major depression.

Methods: Data from 703 individuals with major depression recruited from the research database of the sleep laboratory of the Erasme Hospital were analysed. Only individuals with a diagnosis of type 2 diabetes according to the diagnostic criteria of the American Diabetes Association were included in the diabetes group. Logistic regression analyses were conducted to examine clinical and demographic risk factors of type 2 diabetes in major depression.

Download English Version:

<https://daneshyari.com/en/article/6820079>

Download Persian Version:

<https://daneshyari.com/article/6820079>

[Daneshyari.com](https://daneshyari.com)