



Contents lists available at ScienceDirect

Schizophrenia Research

journal homepage: www.elsevier.com/locate/schres

The efficacy of targeted health agents education to reduce the duration of untreated psychosis in a rural population

Eduardo Padilla^{a,b}, Juan Molina^b, Danielle Kamis^b, Maria Calvo^{a,b}, Lee Stratton^b, Sergio Strejilevich^a, Gabriela Gonzalez Aleman^a, Gonzalo Guerrero^{a,b}, Mercedes Bourdieu^{a,b}, Horacio A. Conesa^a, Javier I. Escobar^{c,d}, Gabriel A. de Erausquin^{a,b,*}

^a Fundación de Lucha contra los Trastornos Neurológicos y Psiquiátricos en Minorías (FULTRA), Argentina

^b Hospital Neuropsiquiátrico Néstor Sequeiros, Ministerio de Salud, Provincia de Jujuy, Argentina

^c Roskamp Laboratory for Brain Development, Modulation and Repair, Morsani College of Medicine, University of South Florida, Tampa, FL, USA

^d Rutgers University School of Medicine, Piscataway, NJ, USA

ARTICLE INFO

Article history:

Received 22 February 2014

Received in revised form 26 October 2014

Accepted 29 October 2014

Available online xxxx

Keywords:

Duration of untreated psychosis

Schizophrenia

Primary care intervention

Schedules of Clinical Assessment in

Neuropsychiatry

ABSTRACT

The duration of untreated psychosis (DUP) is a key determinant in the severity of symptoms in patients with schizophrenia. DUP is a modifiable factor that if reduced can improve patient outcome and treatment response. We sought to decrease DUP in rural Argentina by instituting annual training of local health agents to better identify signs of mental illness and offer earlier intervention. DUP was estimated using Schedules of Clinical Assessment in Neuropsychiatry (SCAN). Ongoing training was correlated with a reduction in DUP. Reducing DUP through better screening can decrease the psychosocial burden of disease and improve the trajectory of psychosis.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

The duration of untreated psychosis (DUP) is the amount of time that elapses between the appearance of psychotic symptoms and the initiation of an adequate pharmacological treatment. Even though the onset of psychotic phenomena is often elusive, a variety of methods have been developed to ascertain it objectively. Albeit with the limitations attendant to its accurate determination, DUP predicts the initial clinical course of schizophrenia. Indeed, longer DUP probably has a neurotoxic effect that may result in reduced cortical thickness (Guo et al., 2013) and cognitive impairment (Rapp et al., 2013). Furthermore, longer DUPs are related with shorter periods of symptom remission, longer hospitalizations, less treatment adherence, increased risk of depression and suicide, substance abuse, and higher costs of treatment (Yung et al., 2007). DUP can be influenced by the insidious onset of symptomatology, the lack of public awareness, and the stigma associated with mental illness. Thus, DUP is a modifiable factor whose reduction can improve patient outcomes and increase the effectiveness of

pharmacological treatment (McGlashan, 1999; Penttilä et al., 2013). Hence, the prolongation of DUP is a serious public health issue with complex socio-cultural and pathophysiological implications. Here we describe the impact of yearly training of primary care health agents (HA) in the province of Jujuy, Argentina, on DUP in all cases of never treated psychosis detected during a 7 years surveillance program as part of the Investigation of Movement Abnormalities and Genetic of Schizophrenia study (IMAGES).

The Ministry of Health of the Province of Jujuy has a primary care infrastructure heavily dependent on the use of HA, who are members of each community, typically with high school level education, familiar with the culture, the language, and the attitudes towards self-care in their environment. For HA assignment, the province is divided in 21 districts covering its three geographical regions, namely, the valley, the jungle, and the highlands. Within each district, HA are assigned to monitor, educate, and facilitate access to care for a small number of households, resulting in complete or near complete coverage of the population. Each household is visited at the least two times per year, and HA are charged with the epidemiological surveillance of communicable diseases, health education for primary prevention, vaccination campaigns, prenatal and pediatric well-care, and of engaging and referral of community members in the health system. Prior to the onset of the study reported here, HA had never received training in mental health.

* Corresponding author at: Morsani College of Medicine, University of South Florida, 3515 E. Fletcher Ave., MDC14, Tampa, FL 33613, USA. Tel.: +1 813 974 4716; fax: +1 813 974 3236.

E-mail address: gdeeraus@health.usf.edu (G.A. de Erausquin).

2. Methods

2.1. Targeted training of HA

Beginning in 2004, we carried out annual training of all HA in the province in three sessions, one for each geographical region (valleys, jungle, and highlands). The province has a population of 672,260 according to the 2010 census, with 373,705 belonging to First Nations. The total number of agents on year 1 of the study was 650 and on year 7, 821. Turnover is minimal, so most of the HA had training in all years, and all of them had at least two trainings. Each training session lasted 3 hours and included lectures and printed materials focused on general knowledge about the signs and symptoms of severe mental illness (psychosis, mood or affective symptoms, disabling anxiety, alcohol and substance abuse), basic knowledge about risk factors, and practical instructions on risk management and referral for specialty care. HA fund of knowledge regarding mental health/illness was evaluated by administration of a brief questionnaire immediately before and after sessions to enhance their engagement in the process, but the primary outcome of the training was considered the number of appropriate referrals to the Mental Health system. Prior to 2004, HA did not receive any training whatsoever on mental health or psychiatric diseases. Assessment of training outcomes was achieved by administration of a brief questionnaire to evaluate HA knowledge of mental illness immediately before and after each session. Of note, the primary purpose of the treatment was not necessarily to increase the HA fund of knowledge regarding mental illness but rather to increase awareness and promote early referrals to specialty care.

2.2. Diagnostic ascertainment and assessment of DUP

All protocols used in this study were approved by the Ethics Committee of the Province of Jujuy and by the Morsani College of Medicine IRB at University of South Florida. As part of the IMAGES study, HA were instructed to notify the Direction of Mental Health upon detection of any cases of severe mental illness. A determination was made on first contact of the acuity of the symptoms, and appropriate actions were taken to ensure the safety of the patient and his/her family (including dispatch of an ambulance for hospitalization if necessary and feasible for the person's location). Once safety was ensured, investigators of the IMAGES study were immediately sent to evaluate the new patient in his/her place of residence (or in the Hospital Nestor Sequeiros if admitted). At this point, two possible routes would be followed. If the person and his/her family were interested in participating in research, they were enrolled in the IMAGES study after written informed consent was obtained. Alternatively, if they were not interested, appropriate care was provided free of cost under the universal coverage provincial system. Assessment consisted of semi-structured interviews with the World Health Organization's SCAN (Schedules of Clinical Assessment in Neuropsychiatry), and if criteria for a diagnosis of schizophrenia were met, neuropsychological assessment, neurological exam, and transcranial ultrasound were collected. Inclusion criteria for the study were a diagnosis of DSM-IV schizophrenia (by two independent SCAN raters), age 18 to 65 years, absence of comorbid neurological or psychiatric disorders or substance abuse, and ability to provide informed consent.

2.3. Schedules of clinical assessment in neuropsychiatry (WHO SCAN)

SCAN consists of a series of phenomenological assessment tools, a glossary of psychopathology, and software tools for administering the interviews, and algorithms to render a diagnosis in either ICD or DSM diagnostic systems (Wing et al., 1990). In a study using trained clinicians and videotaped interviews, the agreement per interviewer with regard to the reference diagnoses ranged from 87% (diagnostic group) to 94% (diagnostic cases), and agreement on the syndrome level

Table 1

Interrater reliability of SCAN Part 1 (somatoform, anxiety, mood, and substance use disorders), SCAN Part 2 (psychotic disorders), and individual sections for hallucinations (section 17), experiences of thought interference and replacement of will (section 18), delusions (section 19), motor and behavioral symptoms (section 22), and speech abnormalities (section 24). Three raters independently score videos of the same interviews ($n = 30$). Cohen's kappa scores are provided.

	Kappa	STD
Scan part 1	0.54	0.17
Scan part 2	0.76	0.28
Scan section 17	1.00	0.00
Scan section 18	0.68	0.30
Scan section 19	0.80	0.45
Scan section 22	0.56	0.38
Scan section 24	0.82	0.21

(without duration and interference criteria of DSM-IV) was excellent (Rijnders et al., 2000). Using the same methodology (blind scoring of videotaped interviews), we found very high inter-rater reliability for part 1, part 2, and the psychosis sections of SCAN (Table 1). SCAN has been field tested in 20 centers in 11 countries, including ours (<http://whoscan.org>).

2.4. Estimation of DUP

SCAN provides semistructured items to record the age of onset of prodromal symptoms, the age of onset of first psychotic symptoms, the age of the first recognition of the illness, and the age at which treatment began. These dates were estimated using direct questioning of the subject, collateral information from family members and any alternate sources of information, such as health or school records when available. We estimated onset of DUP as the time from which psychotic symptoms have occurred. In addition, we completed the Symptom Onset in Schizophrenia Inventory (SOS), which rates the presence and dates of onset of general prodromal, positive, negative, and disorganized symptoms, as well as a clinician, family, and patient global rating of onset of illness (Perkins et al., 2000).




Fig. 1. Average DUP (panel a) and average number of referrals to specialty care (panel b) for years with and without training. Error bars represent SD.

Download English Version:

<https://daneshyari.com/en/article/6824182>

Download Persian Version:

<https://daneshyari.com/article/6824182>

[Daneshyari.com](https://daneshyari.com)