Accepted Manuscript

Title: Comparison of cytokines and prooxidants/antioxidants markers among adults with refractory versus well-controlled epilepsy: a cross-sectional study

Authors: Ozlem Ethemoglu, Halil Ay, İsmail Koyuncu,

Ataman Gönel

PII: S1059-1311(18)30258-9

DOI: https://doi.org/10.1016/j.seizure.2018.06.009

Reference: YSEIZ 3210

To appear in: Seizure

Received date: 23-4-2018 Revised date: 4-6-2018 Accepted date: 11-6-2018

Please cite this article as: Ethemoglu O, Ay H, Koyuncu İ, Gönel A, Comparison of cytokines and prooxidants/antioxidants markers among adults with refractory versus well-controlled epilepsy: a cross-sectional study, *Seizure: European Journal of Epilepsy* (2018), https://doi.org/10.1016/j.seizure.2018.06.009

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Comparison of cytokines and prooxidants/antioxidants markers among adults with

refractory versus well-controlled epilepsy: a cross-sectional study

Runnig Head: Comparison of adiponectin, IL-6 and oxidative stress in epilepsy patients.

Authors:

Ozlem Ethemoglu, Asst.Prof.

Harran University School of Medicine, Department of Neurology, Sanlıurfa, Turkey

Halil Ay, Asst.Prof.

Harran University School of Medicine, Department of Neurology, Sanliurfa, Turkey

İsmail Koyuncu, Asst.Prof.

Harran University School of Medicine, Department of Biochemistry and Clinical

Biochemistry, Sanliurfa, Turkey.

Ataman Gönel, Asst.Prof.

Harran University School of Medicine,

Corresponding Author: Ozlem Ethemoglu, Asst. Prof.

Web address: ozlem_uzunkaya@hotmail.com

Phone: +90 532 6159668 Fax: +90 414 318 31 92

Highlights

LDL-C, TC, IL-6 and oxidative stress levels were increased in refractory epilepsy

patients.

Adiponectin and TAS levels was decreased in refractory epilepsy patients.

Refractory epilepsy patients may be more prone to development of atherosclerosis and

cardiovascular disease.

Download English Version:

https://daneshyari.com/en/article/6829739

Download Persian Version:

https://daneshyari.com/article/6829739

<u>Daneshyari.com</u>