

Accepted Manuscript

Title: Scared or scarred: could 'dissociogenic' lesions predispose to nonepileptic seizures after head trauma?

Authors: Stoyan Popkirov, Alan J. Carson, Jon Stone

PII: S1059-1311(18)30109-2
DOI: <https://doi.org/10.1016/j.seizure.2018.04.009>
Reference: YSEIZ 3164

To appear in: *Seizure*

Received date: 13-2-2018
Revised date: 31-3-2018
Accepted date: 10-4-2018

Please cite this article as: Popkirov Stoyan, Carson Alan J, Stone Jon. Scared or scarred: could 'dissociogenic' lesions predispose to nonepileptic seizures after head trauma?. *SEIZURE: European Journal of Epilepsy* <https://doi.org/10.1016/j.seizure.2018.04.009>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



REVISION 1

Ref: SEIZURE_2018_90

Title:

Scared or scarred: could ‘dissociogenic’ lesions predispose to nonepileptic seizures after head trauma?

Category:

Focused review

Authors:

Stoyan Popkirov^{a*}, Alan J. Carson^{b,c}, Jon Stone^b

Affiliations:

^a Department of Neurology, University Hospital Knappschaftskrankenhaus Bochum, Ruhr-University Bochum, Bochum, Germany

^b Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, United Kingdom

^c Scottish Neurobehavioural Rehabilitation Unit, Royal Edinburgh Hospital, Edinburgh, United Kingdom

*Corresponding author:

Dr. Stoyan Popkirov, Klinik für Neurologie, Universitätsklinikum Knappschaftskrankenhaus Bochum, In der Schornau 23-25, 44892 Bochum, Germany.

Email: popkirov@gmail.com.

Phone: +49 234 299-3701.

Fax: +49 234 299-3719

Highlights

- Traumatic brain injury (TBI) is common in psychogenic nonepileptic seizures (PNES).
- A possible connection is seen in maladaptive symptom modelling.
- Effects of mild brain injury can include disruption of functional connectivity.
- This disruption might temporarily facilitate dissociation and PNES development.
- Mild TBI might be a cause of “dissociogenic” lesions in some patients.

Download English Version:

<https://daneshyari.com/en/article/6829837>

Download Persian Version:

<https://daneshyari.com/article/6829837>

[Daneshyari.com](https://daneshyari.com)