

Accepted Manuscript

Title: The burden of headache in people with epilepsy

Author: R. Mameniškienė I. Karmonaitė R. Zagorskis

PII: S1059-1311(16)30124-8

DOI: <http://dx.doi.org/doi:10.1016/j.seizure.2016.07.018>

Reference: YSEIZ 2771

To appear in: *Seizure*

Received date: 31-3-2016

Revised date: 29-7-2016

Accepted date: 29-7-2016



Please cite this article as: <http://dx.doi.org/>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

1. 83.2% of people with epilepsy have a burden of a headache;
2. Every second person with epilepsy has 3rd or 4th grade of HALT index;
3. The frequency, duration and type of a headache influence the quality of life of PWE;
4. The most common type of headache among people with epilepsy is tension type headache;
5. There was no gender-related difference in the prevalence of migraine in PWE.

Download English Version:

<https://daneshyari.com/en/article/6830368>

Download Persian Version:

<https://daneshyari.com/article/6830368>

[Daneshyari.com](https://daneshyari.com)