Accepted Manuscript

Adrenal Tumor Segmentation Method for MR Images

Mücahid Barstuğan, Rahime Ceylan, Semih Asoglu, Hakan Cebeci, Mustafa Koplay

 PII:
 S0169-2607(18)30853-8

 DOI:
 10.1016/j.cmpb.2018.07.009

 Reference:
 COMM 4755



To appear in: Computer Methods and Programs in Biomedicine

Received date:	7 June 2018
Revised date:	11 July 2018
Accepted date:	16 July 2018

Please cite this article as: Mücahid Barstuğan, Rahime Ceylan, Semih Asoglu, Hakan Cebeci, Mustafa Koplay, Adrenal Tumor Segmentation Method for MR Images, *Computer Methods and Programs in Biomedicine* (2018), doi: 10.1016/j.cmpb.2018.07.009

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- Adrenal tumors can be adherent to liver, spleen, spinal cord, kidney, and this situation prevents an accurate segmentation of adrenal tumors.
- In addition, low contrast, non-standardized shape and size, homogeneity, and heterogeneity of the tumors are considered as problems in segmentation.
- This study proposes a computer-aided diagnosis (CAD) system to segment adrenal tumors by eliminating the above problems.
- The performance of the proposed method was assessed on 113 Magnetic Resonance (MR) images using seven statistical metrics.
- As a result, an efficient framework is presented on segmentation of adrenal tumors for MR images, especially for cyst-based tumors.

Download English Version:

https://daneshyari.com/en/article/6890707

Download Persian Version:

https://daneshyari.com/article/6890707

Daneshyari.com