HOSTED BY

Contents lists available at ScienceDirect

### Egyptian Journal of Basic and Applied Sciences

journal homepage: www.elsevier.com/locate/ejbas



Full Length Article

# Prevalence and characterization of carbapenem-resistant *Klebsiella* pneumoniae isolated from intensive care units of Mansoura University hospitals



Dalia Moemen\*, Doaa T. Masallat

Department of Medical Microbiology and Immunology, Faculty of Medicine, Mansoura University, Egypt

#### 1. Introduction

Klebsiella pneumoniae (K. pneumoniae) is a Gram negative, facultative anaerobic bacilli that has great potential for significant morbidity and mortality in acute care settings, particularly in immunocompromised patients [1]. Moreover, K. pneumoniae is one of the most common organisms showing multiple antibiotic resistance worldwide [2]. These bacteria easily acquire and transfer drug resistance genes through plasmids and transposons [3]. Acquisition of these genes leads to production of β-lactamases of which extended spectrum β-lactamases (ESBLs) are the most common [3]. ESBLs are capable to hydrolyze extended spectrum penicillins, cephalosporins and monobactams, leaving the carbapenem group of  $\beta$ -lactam antibiotics as the only choice for therapy, hence carbapenem antibiotics are used as a last resort to treat infections caused by these multidrug resistant organisms [4,5]. However, there has been emergence of carbapenem resistant Enterobacteriaceae, most commonly carbapenem resistant K. pneumoniae (CRKP), which have a worldwide prevalence [6], due to high antibiotic use, self-medication by patients and lack of implementation of antibiotic policies in hospitals [7]. Mechanisms described for carbapenem resistance include, production of different classes of carbapenemase, hyperproduction of AmpC β-lactamase with an outer membrane porin mutation, and production of ESBL with a porin mutation or drug efflux. Production of carbapenemases is the most commonly reported mechanism of carbapenem resistance in K. pneumoniae [8]. Carbapenemases are β-lactamase enzymes that are capable of hydrolyzing all beta-lactam antibiotics, including monobactams, extended spectrum cephalosporins and carbapenem [9]. The most common carbapenemases include Verona integron metallo-beta-lactamases types (VIM), imipenemase (IMP) types, Klebsiella pneumoniae carbapenemase (KPC), oxacillinase-48 (OXA-48), and New Delhi metallo-betalactamase-1 (NDM-1), encoded by carbapenem resistance determining genes blaVIM, blaIMP, blaKPC, blaOXA-48-like, and blaNDM, respectively [10]. Phenotypic assays are used to identify activity of carbapenemase while molecular assays have developed to identify

E-mail address: dr\_daliamoemen@yahoo.com (D. Moemen).

carbapenemase encoding genes [10,11]. Analysis of hospital surveillance data by the Center for Disease Control and Prevention (CDC) suggested that 8% of all Klebsiella isolates are carbapenem resistant [12]. Other studies showed that it accounted for 5-24% of Klebsiella isolates identified in hospitalized patients [13]. In Egypt, carbapenem resistance is emerging and alarming, one study reported that carbapenem resistance was detected in 44.3% of K. pneumoniae isolates in Suez Canal university hospitals [14]. The detection of carbapenem resistance is essential for the proper choice of antibiotic therapy as well as infection control measures to prevent dissemination of resistant strains in hospital settings. Therefore we set out this study to determine the prevalence of carbapenem resistance and carbapenemase encoding genes among clinical K. pneumoniae isolates obtained from patients at intensive care units (ICUs) of Mansoura university hospitals (MUHs), taking in consideration that carbapenems are frequently used as an empiric therapy in ICUs at our institution.

#### 2. Subjects and methods

#### 2.1. Study design

This was a cross-sectional study, between January 2015 and March 2016. All patients admitted to different ICUs of MUHs and had confirmed infection by *K. pneumoniae*, were enrolled in this study. The ICU bed numbers range from 4 to 27, with a median of 10. This study was conducted with approval from the Medical Research Ethics Committee, Mansoura University.

#### 2.2. Case definition

A patient with a culture positive for *K. pneumoniae* was deemed to have an infection if *K. pneumoniae* was isolated from a sterile site (e.g., blood, peritoneal fluid, cerebrospinal fluid, or pleural fluid) in combination with clinical signs and symptoms of infection. Pneumonia was diagnosed on the basis of clinical signs and symptoms (cough, dyspnea, fever), the appearance of infiltrate on chest radiography and heavy growth of organisms in purulent tracheal secretions or bronchoalveolar lavage fluid (>10<sup>4</sup> colony forming units/ml). These samples were examined after Gram staining for the detection and quantification of leukocytes and organisms.

<sup>\*</sup> Corresponding author at: Mansoura University, Microbiology Department, Faculty of Medicine, Elgomhouria St., Mansoura 35516, Egypt.

A diagnosis of urinary tract infection required the isolation of at least 10000 microorganisms/ml associated with at least two of the following: dysuria; frequency; and/or pyuria (>10 white blood cells per high power field) [15].

#### 2.3. Sample collection and processing

Samples differed according to site of infection; they included blood, peritoneal fluid, urine, respiratory secretions, wound swabs, and cerebrospinal fluid. Only one isolate per patient was included. All collected samples were processed in the Microbiology Diagnostics and Infection Control Unit (MDICU), within 1–2 h of collection.

#### 2.4. Isolation and identification of K. pneumoniae

Bacterial identification to the species level were carried out by colonial morphology on blood and MacConkey's agar plates (oxoid, UK), Gram stained films, biochemical reactions including oxidase, motility, indole, methyl red, voges-proskauer, citrate and urease tests and confirmed with API 20E (BioMerieux, Inc., Hazelwood, MO).

#### 2.5. Antibiotic susceptibility testing

The disk diffusion method was employed to determine the susceptibility profile, carried out on a Muller-Hinton agar, as recommended by CLSI M100-S24 [16]. The following disks, provided by (Oxoid, UK) were used: imipenem (IPM) (10  $\mu$ g), meropenem (MEM) (10  $\mu$ g), ertapenem (ERT) (10  $\mu$ g), amoxicillin/clavulanic acid (AMC) (30  $\mu$ g), piperacillin/tazobactam (TZP) (110  $\mu$ g), ceftazidime (CAZ) (30  $\mu$ g), cefepime (FEP) (30  $\mu$ g), ciprofloxacin (CIP) (10  $\mu$ g), gentamicin (CN) (10  $\mu$ g), amikacin (AK) (30  $\mu$ g). CRKP was defined as *K. pneumoniae* isolates that test intermediate or resistant to ertapenem, according to breakpoints defined by the CLSI [16].

#### 2.6. Detection of carbapenemase production

Carbapenemase production was confirmed by the following tests: a) Modified Hodge test (MHT): a 1:10 dilution of 0.5 McFarland standard suspension of E. coli ATCC 25922 was made (adding 0.5-4.5 ml of saline) and swab streaked all over the plate. Then, 10 µg ertapenem disk was placed in the center of the test area. The test isolate was streaked in a straight line from the disk to the edge of the plate. Interpretation of negative and positive tests was done according to CLSI, (2014) [16]. b) Boronic acid synergy test: was done by streaking 0.5 McFarland standard suspension of the test isolate on a plate. 10 µg ertapenem and 400 µg of phenylboronic Acid (PBA) disks were then placed on the inoculated plate 15 mm apart center to center, and incubated for 24 h. The presence of enhanced growth inhibition zone between the carbapenem disk and boronic acid disk was considered positive for KPC enzyme production [17] c) Ethylene diammine tetra acetic acid (EDTA) test: 0.5 McFarland standard suspension of the test isolate was spread on the surface of a Mueller Hinton Agar plate. Two 10 µg ertapenem disks were placed on the agar 15 mm apart center to center. 10 µl of 0.5 M EDTA was added to one of the ertapenem disk to get 750 µg concentration and incubated at 37 °C overnight. Increase of inhibition zone diameter of more than 5 mm in the disk potentiated with the EDTA was considered positive for metallo- $\beta$ -lactamase production [18].

#### 2.7. Detection of carbapenemase encoding genes

DNA extraction was performed by the boiling method using the CDC protocol [19]. Polymerase chain reaction (PCR) was carried out

in a thermal cycler machine (MJ Research, Inc., USA). The genes blaIMP, blaVIM, blaKPC, blaOXA-48-like and blaNDM-1 were amplified using primers and conditions as described by Karuniawati et al. (2013) [7]. A volume of 1  $\mu L$  of template DNA was added to a final volume of 25  $\mu L$  PCR mixture comprising 12.5  $\mu L$  of Taq PCR Master Mix (Fermentas, UK), including, 1× PCR buffer, 1.5 mmol/L MgCl2, 0.15 mmol/L dNTP, and 1.25 IU Taq DNA polymerase, 1  $\mu L$  of 0.8  $\mu$ mol/L each primer (except OXA 0.4  $\mu$ l) and 9.5  $\mu L$  of sterile distilled water. The primer pairs were tested in simplex PCR (only one gene screened, for blaNDM-1 gene) and with a multiplex approach. The amplicons were analyzed by electrophoresis in a 1.5% agarose gel.

#### 2.8. Data analysis

Data analysis was done in STATA version 12. Out comes were presented as proportions and percentages in a tabular form.

#### 3. Results

#### 3.1. Patients and isolates characteristics

This study included 125 patient with confirmed *K. pneumoniae* infections, among which 42 had CRKP infections. CRKP isolates were obtained from 23 male and 19 female patients. Patients were generally elderly (median age was 60 years, with a range of 43–77 years). Respiratory samples (62%) was the predominant source of CRKP, followed by urine (14%), wound (9.5%), blood (9.5%), and catheter tip (5%) samples.

#### 3.2. Susceptibility to antibiotics

Using current breakpoints recommended by CLSI (M100-S24) for carbapenem interpretation, 42 out of 125 (33.6%) *K. pneumoniae* isolates were nonsusceptible (intermediate and resistant) to ertapenem. 25.6% and 20% were nonsusceptible to meropenem and imipenem respectively. The full antibiogram for the tested drugs is seen in Table 1. Two isolates were resistant to all tested antibiotics (panresistant), however these isolates clinically respond well to the last-resort antimicrobials colistin and tigecycline.

#### 3.3. Prevalence of carbapenemase activity based on phenotypic tests

Carbapenamase activity was detected in 26/42 (61.9%) by MHT method, 22/42 (52.4%) by Boronic acid screen and 5/42 (11.9%) by the EDTA test. 18 of the isolates were positive for both the MHT and Boronic acid methods. Overall, 35 (83.3%) of the 42 CRKP were positive for the production of one or more carbapenemases. Details of the carbapenemase activity among the isolates are shown in Table 2.

#### 3.4. Prevalence and distribution of carbapenemase genes

Based on the PCR assays, 39/42 of the CRKP isolates were positive for one or more carbapenemase genes, while none of the carbapenemase genes tested (blaKPC, blaNDM, blaVIM-1, blaIMP, and blaOXA-48-like) were detected in three isolates tested. Furthermore, these three isolates did not phenotypically express a carbapenemase, according to the MHT, Boronic acid and EDTA tests. Carbapenem resistance of these isolates is likely due to a combination of ESBLs and changes in outer membrane proteins (ESBL/Omp) [20]. Five of the 39 (12.8%) carbapenemase gene carrying isolates harbored two or more genes. As shown in Table 3, the most prevalent gene was *bla*KPC 47.8% followed by *bla*VIM-1

#### Download English Version:

## https://daneshyari.com/en/article/6952390

Download Persian Version:

https://daneshyari.com/article/6952390

<u>Daneshyari.com</u>