

## Accepted Manuscript

Title: Efficient degradation of imipramine by iron oxychloride-activated peroxymonosulfate process

Authors: Yiqun Chen, Yanpeng Liu, Li Zhang, Pengchao Xie, Zongping Wang, Aijiao Zhou, Zheng Fang, Jun Ma



PII: S0304-3894(18)30211-5  
DOI: <https://doi.org/10.1016/j.jhazmat.2018.03.050>  
Reference: HAZMAT 19270

To appear in: *Journal of Hazardous Materials*

Received date: 19-10-2017  
Revised date: 12-3-2018  
Accepted date: 26-3-2018

Please cite this article as: Chen Y, Liu Y, Zhang L, Xie P, Wang Z, Zhou A, Fang Z, Ma J, Efficient degradation of imipramine by iron oxychloride-activated peroxymonosulfate process, *Journal of Hazardous Materials* (2010), <https://doi.org/10.1016/j.jhazmat.2018.03.050>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# Efficient degradation of imipramine by iron oxychloride-activated peroxymonosulfate process

Yiqun Chen<sup>a</sup>, Yanpeng Liu<sup>b,c</sup>, Li Zhang<sup>b,c</sup>, Pengchao Xie<sup>b,c,\*</sup>, Zongping Wang<sup>b,c</sup>, Aijiao Zhou<sup>b,c,\*</sup>,

Zheng Fang<sup>a</sup>, and Jun Ma<sup>d</sup>


<sup>a</sup> School of Civil Engineering, Wuhan University, Wuhan 430072, China

<sup>b</sup> School of Environmental Science and Engineering, Huazhong University of Science and Technology (HUST), Wuhan 430074, China

<sup>c</sup> Key Laboratory of Water & Wastewater Treatment (HUST), MOHURD, Wuhan 430074, China

<sup>d</sup> State Key Laboratory of Urban Water Resource & Environment, Harbin Institute of Technology, Harbin 150090, China

## Graphic Abstract



## Highlights

- Synthesized FeOCl was firstly used to activate PMS.
- Sulfate and hydroxyl radicals were identified as reactive species in FeOCl/PMS.
- PMS was mainly activated by the unsaturated iron atoms and oxo-bridged configuration.

\* Corresponding authors.

E-mail address: [pengchao.x@gmail.com](mailto:pengchao.x@gmail.com) (P. Xie); [ajzhou@hust.edu.cn](mailto:ajzhou@hust.edu.cn) (A. Zhou)

Download English Version:

<https://daneshyari.com/en/article/6968361>

Download Persian Version:

<https://daneshyari.com/article/6968361>

[Daneshyari.com](https://daneshyari.com)