

Accepted Manuscript

Title: Can adsorbent of layered chalcogenide be regenerated?
A case study of norfloxacin adsorbed by layered chalcogenide
in water

Authors: Jian-Rong Li, Lei Xu, Ming-Lai Fu, Huachun Lan



PII: S0927-7757(17)30928-7
DOI: <https://doi.org/10.1016/j.colsurfa.2017.10.029>
Reference: COLSUA 21987

To appear in: *Colloids and Surfaces A: Physicochem. Eng. Aspects*

Received date: 16-7-2017
Revised date: 9-10-2017
Accepted date: 13-10-2017

Please cite this article as: Jian-Rong Li, Lei Xu, Ming-Lai Fu, Huachun Lan, Can adsorbent of layered chalcogenide be regenerated? A case study of norfloxacin adsorbed by layered chalcogenide in water, *Colloids and Surfaces A: Physicochemical and Engineering Aspects* <https://doi.org/10.1016/j.colsurfa.2017.10.029>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Can adsorbent of layered chalcogenide be regenerated? A case study of norfloxacin adsorbed by layered chalcogenide in water

Jian-Rong Li^{a,b}, Lei Xu^a, Ming-Lai Fu^{a,*}, Huachun Lan^{c,d,*}

a. Key Laboratory of Urban Pollutant Conversion, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen, Fujian 361021, P.R. China.

b. Ningbo Key Laboratory of Urban Environmental Processes and Pollution Control, Ningbo Urban Environment Observation and Research Station, Chinese Academy of Sciences, Ningbo, Zhejiang 315800, P.R. China.

c. School of Environment, State Key Joint Laboratory of Environment Simulation and Pollution Control, Tsinghua University, Beijing, 100084, P.R. China.

d. Research Center for Water Purification and Water Ecological Restoration, Tsinghua University, Beijing, 100084, P.R. China.

Graphic Abstract



The cationic norfloxacin was removed by KMS-1 with cation exchange at pH of 4, and it can be regenerated by 1.0 M KCl.

*Corresponding author: Tel.: +865926190762; Fax: +865926190977; E-mail: mlfu@iue.ac.cn; hclan@tsinghua.edu.cn.

Download English Version:

<https://daneshyari.com/en/article/6978028>

Download Persian Version:

<https://daneshyari.com/article/6978028>

[Daneshyari.com](https://daneshyari.com)