Accepted Manuscript



Title: A novel ultradeformable liposomes of Naringin for anti-inflammatory therapy

Authors: María Pleguezuelos-Villa, Silvia Mir-Palomo, Octavio Díez-Sales, M.A. Ofelia Vila Buso, Amparo Ruiz Sauri, Amparo Nácher

PII: S0927-7765(17)30833-0
DOI: https://doi.org/10.1016/j.colsurfb.2017.11.068
Reference: COLSUB 9019

To appear in: *Colloids and Surfaces B: Biointerfaces*

Received date: 28-9-2017
Revised date: 25-11-2017
Accepted date: 30-11-2017

Please cite this article as: María Pleguezuelos-Villa, Silvia Mir-Palomo, Octavio Díez-Sales, M.A.Ofelia Vila Buso, Amparo Ruiz Sauri, Amparo Nácher, A novel ultradeformable liposomes of Naringin for anti-inflammatory therapy, Colloids and Surfaces B: Biointerfaces https://doi.org/10.1016/j.colsurfb.2017.11.068

A novel ultradeformable liposomes of Naringin for anti-inflammatory therapy

María Pleguezuelos-Villa[1, *], Silvia Mir-Palomo[1], Octavio Díez-Sales [1,2], M.A. Ofelia Vila Buso[3], Amparo Ruiz Sauri [4], Amparo Nácher[1, 2].

[1]Department of Pharmacy, Pharmaceutical Technology and Parasitology, Faculty of Pharmacy, University of Valencia, Av. Vicent Andrés Estellés s/n, 46100, Burjassot, Valencia, Spain.

[2]Instituto Interuniversitario de Investigación de Reconocimiento Molecular y Desarrollo Tecnológico (IDM), Universitat Politècnica de València, Universitat de València, Av. Vicent Andrés Estellés s/n, 46100, Burjassot, Valencia, Spain.

[3]Department of Physical Chemistry, Faculty of Pharmacy, University of Valencia, Av. Vicent Andrés Estellés s/n, 46100, Burjassot, Valencia, Spain.

[4] Department of Pathology, University of Valencia, Av. Blasco Ibañez 17, 46010 Valencia, Spain.

***Corresponding author**: M. Pleguezuelos-Villa; e-mail address: maplevi@alumni.uv.es

The article has 4636 words including ackwonledgment and references. The abstract has 127 words. There are 5 figures and 2 tables.

Highlights

- Nanovesicles were small in size and monodispersed independently of the NA concentration.
- No citotoxity effects were observed in 3T3 fibroblasts treated with ultradeformable liposomes.