



Synthesis of S-licarbazepine by asymmetric reduction of oxcarbazepine with *Saccharomyces cerevisiae* CGMCC No. 2266

Zhi-Min Ou^{a,*}, Han-Bing Shi^b, Xing-Yuan Sun^b, Wen-He Shen^c

^a Pharmaceuticals College, Zhejiang University of Technology, Hangzhou, Zhejiang 310014, China

^b The Third Affiliated Hospital of Qiqihar Medical College, Qiqihar, Heilongjiang 161000, China

^c Zhejiang Jiuzhou Pharma Science & technology Co., Ltd., Hangzhou, Zhejiang 310051, China

ARTICLE INFO

Article history:

Received 6 May 2011

Received in revised form 18 June 2011

Accepted 4 July 2011

Available online 13 July 2011

Keywords:

S-licarbazepine

Asymmetric reduction

Oxcarbazepine

Biotransformation

ABSTRACT

S-licarbazepine was synthesized by asymmetric reduction of oxcarbazepine with CGMCC No. 2266. The optimum batch reduction conditions were found to consist of a reaction time of 36 h, temperature of 30 °C, and initial pH value of 7.0. The optimum concentration of the glucose co-substrate was found to be 0.3 mol L⁻¹. The addition of glucose contributed to *in situ* regeneration of NADPH in cells and improved conversion. Conversion increased with the addition of more biomass and with a decrease in the initial concentration of substrate. Within the membrane reactor, a continuous reduction process was used to improve production efficiency and reduce the inhibition of high-concentration substrate upon reduction. The optimum flux was found to be 20 ml h⁻¹. S-licarbazepine yield was 3.7678 mmol L⁻¹ d⁻¹ in continuous reduction over four days. The enantiometric excess of S-licarbazepine was 100% for both batch and continuous reduction processes.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

S-licarbazepine (S-LC) can be used in synthesis of eslicarbazepine acetate by esterification with acetic acid. Eslicarbazepine acetate [S-(–)-10-acetoxy-10,11-dihydro-5H-dibenz[–b,f]azepine-5-car-Boxamide, ESL], formerly known as BIA 2-093, is a novel central nervous system (CNS)-active compound with anticonvulsant activity [1–5]. It behaves as a voltage-gated sodium channel blocker and is currently under clinical development for the treatment of epilepsy and bipolar disorder [6–8]. In humans, ESL is more rapidly reduced by liver esterases to the major active metabolite S-licarbazepine than oxcarbazepine (OXC), a new antiepileptic drug approved for the treatment of partial onset seizures and generalized tonic-clonic seizures [9–18].

This paper discusses in detail the synthesis of S-licarbazepine by asymmetric reduction of oxcarbazepine with CGMCC No. 2266 as a catalyst. This is the first report on biosynthesis of S-licarbazepine using microorganism catalysts. Industry has adopted chemical synthesis as a preferred method of production of S-licarbazepine. No references exist on the enzymatic asymmetric reduction of OXC to S-LC. S-licarbazepine can be obtained by asymmetric reduction of oxcarbazepine within microorganisms (Fig. 1). Chiral alcohol, an intermediate of many different pharmaceuticals, can be obtained

by asymmetric reduction of prochiral ketones through biotransformation [19–22]. Asymmetric reduction with microorganism catalysts has many advantages for the production of chiral compounds. These include safety and reliability, low cost, scalability, a lack of environmentally dangerous by-products, the fact that cell culture facilities already exist in many laboratories, and the purity of the final chiral product. In addition, the regeneration of coenzyme (NADH and NADPH) *in situ* can improve conversion because of many kinds of enzyme exist in cells [23,24].

2. Experimental procedure

2.1. Reagents and instruments

Oxcarbazepine and S-licarbazepine were purchased from Zhejiang Jiuzhou Pharma Science & Technology Co., Ltd. (China). Agilent HPLC 1200 equipped with Daicel OD-H chiral column from Daicel Company was used in detection of substrate and product content and enantiometric excess of S-licarbazepine. The membrane reactor and ultrafiltration membrane (MWCO 30 kDa) were from Shanghai Mosu Company (China).

2.2. Microorganism cultivation

Saccharomyces cerevisiae CGMCC No. 2266 was obtained from the soil in the vicinity of Hang Zhou West Lake Brewery and preserved in China General Microbiology Culture Collection Cen-

* Corresponding author. Tel.: +86 0571 88320320; fax: +86 0571 88320320.
E-mail address: oozzmm@163.com (Z.-M. Ou).




Fig. 1. Synthesis of ESL by asymmetric reduction of OXC.

ter on November 26, 2007. Slant medium for strain storage was composed of 10 g L^{-1} malt juice, 3 g L^{-1} yeast extract, 5 g L^{-1} peptone, 10 g L^{-1} glucose and 20 g L^{-1} agar. The liquid yeast culture medium was composed of 30 g L^{-1} glucose, 3 g L^{-1} yeast extract, 5 g L^{-1} $(\text{NH}_4)_2\text{SO}_4$, 0.25 g L^{-1} MgSO_4 , 1 g L^{-1} $\text{K}_2\text{HPO}_4 \cdot 3\text{H}_2\text{O}$, and 1 g L^{-1} KH_2PO_4 . The strain picked from slant medium was inoculated into 100 ml liquid culture medium and cultivated in 30°C shaker (200 r min^{-1}) for 24 h. Then 10 ml of the cell suspension was inoculated into 100 ml liquid medium. After 24 h of cultivation, the cells were harvested and used in reduction.

2.3. Preparation of oxcarbazepine alcohol solution

Oxcarbazepine alcohol solution was prepared by the addition of oxcarbazepine to hot alcohol, which was then cooled to room temperature. The concentration of the oxcarbazepine alcohol solutions were 0.793, 1.982, 3.964, 5.987, and $7.928 \text{ mmol L}^{-1}$.

2.4. Batch reduction

The cultivated cells were separated from the culture medium by centrifugation. The cells from the sediment were washed twice with sterile water and separated by centrifugation before being in reduction. The sediment obtained above and 10 ml oxcarbazepine alcohol solution were added to a flask containing 100 ml of 0.2 mol L^{-1} phosphate buffered solution for reduction. The reaction was stopped by centrifugation at the end of the reduction period. The supernatant obtained above was extracted by ethyl acetate. The ethyl acetate layer was detected for determination of the substrate and product content by Agilent HPLC 1200.

2.5. Continuous reduction

Fig. 2 shows the continuous reduction of oxcarbazepine in the membrane reactor. One hundred milliliters of oxcarbazepine alcohol solution was pumped from vessel 1 into a continuously operated stirred membrane reactor (reactor 2). The flow rate of oxcarbazepine alcohol solution was $10\text{--}50 \text{ ml h}^{-1}$. The 200 ml of reduction mixture in reactor 2 was composed of cells, 0.3 mol L^{-1} glucose, and phosphate buffer (pH 8.0). The cell dry weight in the mixture was 28 g . The cells were trapped by an ultrafiltration membrane (MWCO 30 kDa). The membrane was precoated with 1 mg of bovine serum albumin per ml reactor volume to prevent cell adsorption. The whole reactor was sterilized with 0.01% peracetic




Fig. 2. Continuous reduction process. (1) Substrate alcohol solution storage tank; (2) membrane reactor for continuous reduction; (3) ultrafiltration membrane (MWCO 30 kDa); and (4) peristaltic pump.

acid before use. Percolating fluid from reactor 2 was pumped into vessel 1 for cyclic utilization. Reaction volume, reaction temperature (28°C), and stirring speed (160 r min^{-1}) remained unchanged. The flux pumped into vessel 1 was equal to that pumped out of vessel 1. At the end of reduction, the liquid in vessel 1 was extracted by ethyl acetate and the ethyl acetate layer was detected for determination of the substrate and product content by Agilent HPLC 1200.

2.6. Production of S-licarbazepine by continuous reduction

After $7.928 \text{ mmol L}^{-1}$ of OXC dissolved in 100 ml alcohol solution was completely converted to S-LC, another 100 ml alcohol solution containing $7.928 \text{ mmol L}^{-1}$ of OXC was pumped into vessel 2 for continuous reduction. The flux was 20 ml h^{-1} . The process was repeated until the cells in vessel 2 lost reduction activity.

3. Results and discussion

3.1. Influence of initial substrate concentration and reaction time on batch reduction

Fig. 3 shows that conversion increased gradually with reaction time and the rate of increase was very minor after 36 h. The optimum reaction time is therefore 36 h. Conversion decreased with as the concentration of initial substrate increased. This was probably because the reduction activity of cell was inhibited by high con-




Fig. 3. Influence of initial substrate concentration and reaction time on conversion 30°C , initial pH value of reaction mixture 7.0, 160 r min^{-1} , cell concentration 140 g L^{-1} .

Download English Version:

<https://daneshyari.com/en/article/70197>

Download Persian Version:

<https://daneshyari.com/article/70197>

[Daneshyari.com](https://daneshyari.com)