



ELSEVIER

Available online at www.sciencedirect.com

ScienceDirect

<http://www.elsevier.com/locate/biombioe>

Short Communication

Allocation of five macroelements and quality of fuels derived from Norway spruce wood obtained by thinning operations

Leonardo Cerasino ^a, Nicola La Porta ^{a,b,*}

^a Sustainable Agro-Ecosystems and Bioresources Department, IASMA Research and Innovation Centre, Fondazione Edmund Mach, Via E. Mach 1, 38010, San Michele all'Adige, Trento, Italy

^b MOUNTFOR Project Centre, European Forest Institute, Via E. Mach 1, San Michele all'Adige, Trento, Italy

ARTICLE INFO

Article history:

Received 14 June 2013

Received in revised form

4 August 2014

Accepted 13 August 2014

Available online xxx

Keywords:

Wood minerals

Ecological sustainability

Soil fertility

Tree harvesting systems

Tree mass allocation

Fuel chip quality

ABSTRACT

The use of forest biomass for energy production is growing in Europe and biomass energy plants market is constantly increasing. However, there is the need to define the environmental sustainability issues dealing with the emerging renewable energy scenario. In particular, the polluting emissions (i.e. PM_x, NO_x and ozone) caused by the biomass combustion heavily impact on the air quality. In this context, the elemental characterization of the wood and the element allocation in the different tree organs, can provide important information about the quality of the derived wood fuels and give hints about the choice of the most appropriate combustion technique and/or the right wood fuel for a given combustion technique. Moreover, since elements have different concentrations in the different plant tissues, the preventive knowledge of the elements allocation can lead to the identification of the best harvesting strategy aimed at producing wood fuel with the lowest possible ash forming elements and environmental impact.

This work focuses on the allocation in three tree compartments (foliage, branches and stem) of five important macroelements (K, Mg, Ca, N and P) in Norway spruce (*Picea abies*), and points out the possible effects of different harvesting strategies and tree age on the quality of the wood fuels. Results suggest that the Stem Only Harvesting is preferable to Whole Tree Harvesting system in terms of prevention of mineral content loss, as well as is preferable to avoid forest biomass from young trees because of the poorer fuel quality of the wood chips.

© 2014 Elsevier Ltd. All rights reserved.

1. Introduction

Today there is an increased activity on use of wood as producer of energy with main drivers being climate change,

shortage and increasing prices of fossil fuel sources and safety in energy supply.

However, intensive use of forest biomass is debated since fundamental ecological processes may be influenced

* Corresponding author. Fondazione Edmund Mach, Via E. Mach 1, 38010, San Michele all'Adige, Trento, Italy. Tel.: +39 0461 615 396, +39 338 7888736 (mobile); fax: +39 0461 650 956.

E-mail address: nicola.laporta@fmach.it (N. La Porta).

<http://dx.doi.org/10.1016/j.biombioe.2014.08.006>

0961-9534/© 2014 Elsevier Ltd. All rights reserved.

negatively thus making up a trade-off with the benefits of using an otherwise sustainable source of energy [1].

The use of “wood fuels” presents several drawbacks from environmental and technological points of view: during combustion and biomass conversion processes, many undesired phenomena may happen, like corrosion, formation of deposits, production of ash, harmful gases (such as NO_x , HCl) and particulate matter (PM_x). These negative phenomena are linked to the presence of some elements during combustion, especially those indicated as “ash forming elements” (Si, Al, Fe, Ca, Mg, Na, K, P, S, Cl) [2–5]. Therefore, the quality of a given biomass is positively correlated with the energy content (linked to the presence of C, H and O), but is negatively correlated to the presence of these elements. Ash elements in particular are very variable as they depend on the plant species, the growing site, the plant physiology (season, age, assortment) and the plant management [6]. The knowledge of the relationships between all these and other factors [7] and the elemental content in the biomass could provide valuable criteria for sustainable management of wood stocks for energy purposes [8].

In this work we have focused our attention to the Norway spruce (*Picea abies*) species, which is one of the most important wood fuel sources in Europe. We investigated the effect of the plant age upon the allocation of five important elements, i.e. four ash elements (K, Mg, Ca, and P) plus N, which is endowed with the emission of harmful gases. We then used these data to determine which harvesting strategy, between Stem Only Harvesting (SOH) and Whole Tree Harvesting (WTH), could be the most suitable for getting a better quality biomass and for being the more ecologically sustainable.

2. Material and methods

2.1. Experimental sampling

Sampling was carried out in two different stands representative of the Trentino (Eastern Italian Alps) forests. Stand 1 was a commercial thinning plot (middle aged stands) located in the Commune of Molina di Fiemme (46°16' N, 11°26' E) at elevation ca. 1020 m a.s.l., with Norway spruce 87%, slope gradient 35%, soil type Brown Podzol and humus type Moder. Stand 2 was an unthinned precommercial plot (young stands) located in the commune of Ziano (46°17' N, 11°34' E) at elevation ca. 1080 m a.s.l., with Norway spruce 90%, slope gradient 29%, soil type Brown Podzol and humus type Mull. Both soil stands were classified as medium fertility by the Forest Management Plan of the Autonomous Province of Trento [9]. Collection of biomass samples [7] was carried out in September 2011 and in each stand, four representative trees were randomly sampled. The stands were fully stocked, with a closed canopy and no visual crown damage. Diameters at breast height (DBH) were measured for all trees. Young stand was ca. 25 years old and averaged 14.8 cm DBH while middle aged stand was ca. 64 years old and averaged 32.4 cm DBH. Care was taken to select trees with regular growth and absence of tension and compression wood, or any decay, which can result in natural anomalous concentrations of certain elements. Wood core samples, including bark, were taken at DBH from the base of the trunk, using a chainsaw to extract ca. 20 cm thick radial

cross sections. These sections were double wrapped into clear contamination-free polyethylene plastic bags and kept chilled to limit sap migration before arrival in the laboratory. After arrival in the laboratory of the Fondazione Mach, wood samples were chipped into pieces of less than 2 cm. Living branch and needle samples were collected from the upper part of the crown on the same trees and treated as a bulk sample. Needles were free of fungal damages, herbivores, or decay and characterized by good growth, without discoloration or defoliation. Needles of all ages were manually separate and washed with deionised water. Needle, branch and stem wood samples, including bark, were dried in oven to constant weight at 40 °C. For each tree ca. 8 kg of dry core wood, 8 kg of dry branches and 1 kg of dry needles were collected, from which 200 g was grounded and homogenised and from which 100 mg were further chemically analysed for nutrient elements.

After homogenization, four subsamples of each dried grounded material were processed for chemical analysis. A total of 24 plant samples were analysed in four replicates (96 total analyses).

2.2. Chemical analysis

Ultra-pure grade reagents were used. Approx. 100 mg of dried grounded plant material were digested with sulphuric acid, salicylic acid, oxygen peroxide according to standard procedures [10]. The resulting digested solution was adequately diluted with ultra pure water and then the analysis of cations (K, Mg and Ca) was performed by ion chromatography [11], while the analysis of N and P were performed by spectrophotometric methods, measuring nitrate and orthophosphate according to the method of Chapman and Pratt [12]. Limits of quantification were 0.1 g kg⁻¹ dry weight for K and Mg, 1 g kg⁻¹ for Ca, 0.02 g kg⁻¹ for N and P.

2.3. Equipment

The ion chromatographer was a Dionex DX120 (Dionex Corp., Sunnyvale, CA, USA) equipped with a cation exchange column. The spectrophotometer was a Varian CaryBio 50 (Varian Inc., Palo Alto, CA, USA).

3. Results and discussion

Main elemental concentrations found in plant tissues, as mean value and standard error, are reported in Table 1. Tissues have been grouped in three main compartments (stem with bark, branches and foliage). Values are in agreement with those reported in previous studies [13,14] for the same species. It is possible to observe a clear increasing trend in element concentrations from the stem to the foliage. This observation is related to the physiology of the different tissues and can be assumed as valid for all elements in general. In particular, we can note how concentrations in leaves are several times higher than in the other compartments, as leaves are particularly rich in water.

Moreover, element concentrations are generally higher in young than in middle-aged plants in all compartments. P represents a relative exception, as its values are more similar in the two ages. This observation is particularly evident in

Download English Version:

<https://daneshyari.com/en/article/7064257>

Download Persian Version:

<https://daneshyari.com/article/7064257>

[Daneshyari.com](https://daneshyari.com)