



Biooxidation of 1,8-cineole by *Aspergillus terreus*

Carlos García, Paula Rodríguez, Eduardo Días, Horacio Heinzen, Pilar Menéndez*

Cátedra de Farmacognosia y Productos naturales, Laboratorio de Biotatálisis y Biotransformaciones, Departamento de Química Orgánica, Facultad de Química, Universidad de la República (UdelAR), 11800 Montevideo, Uruguay

ARTICLE INFO

Article history:

Received 14 July 2008

Received in revised form 11 February 2009

Accepted 16 February 2009

Available online 28 February 2009

Keywords:

Aspergillus terreus

1,8-cineole

Biooxidation

Hydroxycineole

Monooxygenase

ABSTRACT

Biotransformation of 1,8-cineole by a strain of *Aspergillus terreus* isolated from Eucalyptus leaves was investigated.

This strain produced four oxygenated compounds identified as 2-*exo*-hydroxy-1,8-cineole, 2-*endo*-hydroxy-1,8-cineole, 3-*exo*-hydroxy-1,8-cineole and 3-*endo*-hydroxy-1,8-cineole, with good bioconversion percentage and high stereoselectivity.

© 2009 Elsevier B.V. All rights reserved.

1. Introduction

The monoterpene cyclic ether, 1,3,3-trimethyl-oxabicyclo [2.2.2]octane commonly known as eucalyptol, 1,8-cineole, or cineole (**1**) (see [Scheme 1](#)) is an abundant monoterpene present in the essential oils of several species being the main component of *Eucalyptus globulus* Labill oils [[1](#)].

This monoterpene has been shown to have biological activity against plants [[2–4](#)] microorganisms [[5](#)] and insects [[6,7](#)].

It is extensively used in pharmaceutical preparations for external application and also as a nasal spray [[8](#)]. In turn, due to its pleasant and distinctive flavor, it is also used in the food and cosmetic industries [[9](#)].

The 1,8-cineole, is a cheap raw material and it is easily obtained from *Eucalyptus* essential oils. Its transformation into more valuable compounds is recognized as being of great economic potential to the food, perfume and pharmaceutical industries.

Some studies have been carried out on the oxygenation of 1,8-cineole by chemical [[10,11](#)] and biological processes [[12–19](#)]. However, none of these biotransformations represent a significant breakthrough on the transformation of this abundant and cheap substance into more valuable products.

Obtaining oxygenated derivatives of 1,8-cineole involves stereospecific introduction of molecular oxygen in non-activated carbon atoms, which is not easy to carry out by classical chemical synthesis

[[14](#)]. Besides, the oxidation of this compound by traditional chemical methods usually gives a mixture of stereoisomers. Because of the correlation between structure and biological activity, it is desirable to have methodologies that yield the product in its optically pure form.

The production of new compounds via a biotechnological route offers a number of advantages. One important attribute of microbial biocatalysis is the ability to synthesize products that can be labeled as natural, if derived from natural substrates, and added to foods without being considered as additives [[20](#)].

Our group has been working in the characterization of useful biocatalysts for the synthesis of oxygenated monoterpene derivatives.

In this paper we describe a biocatalytic procedure for the transformation of 1,8-cineole using whole cells of *Aspergillus terreus* isolated from *Eucalyptus* leaves.

This fungus biotransformed 1,8-cineole (**1**) into four oxygenated species (**2**, **3**, **4**, **5**) with good bioconversion percentage and high stereoselectivity ([Scheme 2](#)).

2. Experimental

2.1. Chemical and reagents


1,8-Cineole (99.8%) was provided by the Agroindustrial Technology Center (Cochabamba, Bolivia).

α -Terpineol (90%) was purchased from Aldrich.

m-Chloroperoxybenzoic acid was purchased Sigma–Aldrich.

Ketoconazole and Metronidazole were provided by Roemmers (Montevideo, Uruguay).

* Corresponding author. Tel.: +598 2924 40 68/2929 01 06; fax: +598 2924 1906.
E-mail address: menendez@fq.edu.uy (P. Menéndez).



Scheme 1. (1a) 1,3,3-Trimethyl-2-oxabicyclo[2.2.2]octane and (1b) 1,8-cineole.

2.2. Fungi isolation and identification

Fragments of *Eucalyptus* leaves were transferred to minimal medium BG-11 [21] supplemented with 1,8-cineole (0.5%) as carbon and energy source. After several rounds of serial culturing (1:10 dilutions) at 37 °C and 150 rpm (orbital shaker Sanyo IOXX400.XX2.C), the cell culture was spread onto the minimal medium plates (1.5% agar plates) with 1,8-cineole in the lid, recovering an isolated strain.

For identification of this strain, a culture was grown on Czapek yeast extract agar (CYA) at 5, 25 and 37 °C; malt extract agar (MEA) and 25% glycerol nitrate agar (G25N) at 25 °C. All plates were incubated for 7 days. Fungal identification was done according to Pitt [22]. This strain was kept at the Collection of the Microbiology Department (School of Chemistry, Montevideo, Uruguay) as BFQU 121.


For routine procedures, the strain was grown in PDA slants (DIFCO, Detroit, USA) at 28 °C until sporulation, and then kept at 4 °C.

2.3. Biotransformation assays

YMPG (yeast extract 0.5%, malt extract 1%, bacteriological peptone 0.5%, glucose 1%) was used as the culture media. A spore suspension in sterile physiological serum was used as inoculum reaching a final concentration of 10^6 spores/mL in the culture media.

1,8-Cineole was added after 48 h of incubation such as to attain a final concentration of 0.1% in the culture media.

Biotransformations in the presence of inhibitors were carried out in the same conditions described above. The ketoconazole or metronidazole were added after 48 h of incubation to a final 50 mM [23] concentration in the culture media and 1,8-cineole was added 15 min after the addition of the inhibitor.



Scheme 2. Oxygenated products derived from 1,8-cineole: (2) 2-*exo*-hydroxy-1,8-cineole; (3) 2-*endo*-hydroxy-1,8-cineole; (4) 3-*exo*-hydroxy-1,8-cineole; (5) 3-*endo*-hydroxy-1,8-cineole.

Samples were taken for analysis at 4, 7, 10, and 14 days.

The growth of the fungus, with and without inhibitor, was monitored by measuring dry weight at 100 °C.

Three negative controls were performed, one of them using 1,8-cineole in culture media (without inoculum), a second one inoculated but containing neither substrate nor inhibitor, and a third one similar to the later but including the corresponding inhibitor.

All experiments were carried out in triplicate in 500 mL conical flasks containing 100 mL of culture media each one.

2.4. Extraction and identification of bioconversion products

In order to isolated enough amount of each product to achieve spectral experiments, we carried out the biotransformation experiments by fermentation in 6 conical flasks (2 L) containing 400 mL of culture media each one with 0.1% of 1,8-cineole (total starting material 2400 mg), placed in an orbital shaker Sanyo IOX400 XX2.C, agitation 100 rpm at temperature 28 °C.

The liquid medium was separated from the mycelia by filtration, and then was extracted with CH_2Cl_2 . The mycelia were washed several times with the same solvent. Organic phases were combined and then dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain 1392 mg of the crude mixture of 1,8-cineole **1** and hydroxycineoles **2**, **3**, **4**, and **5**.

Crude mixture reaction was flash-chromatographed on 150 g silica gel SAI (32–63 μm particle size, 60 Å pore size, 40 cm length \times 6.5 cm diameter) using *n*-hexane/ethyl acetate gradient from 4:1 to 2:1.

Each one of these compounds was identified by GC–MS and/or ^1H and ^{13}C NMR.

2.5. Analysis conditions

HRGC analyses were performed in a Shimadzu GC14B equipped with FID and EZ Chrom integration software for data processing. A fused silica capillary column (30 m \times 0.32 mm i.d.) with bonded SE52 (0.40–0.45 μm thickness) was used. Temperature program: 60 °C, 8 min; 60–210 °C at 3 °C/min; injector temperature: 240 °C; detector temperature: 250 °C. Carrier gas: N_2 at 0.50 kg/cm 2 ; injection system: split ratio 1:100.

HRGC–MS was carried out in Shimadzu QP 5500 in the conditions described above, using He as carrier gas. Ionization voltage 70 eV, temperature interface: 250 °C.

The optical purity was determined in a GC–GC Shimadzu GC 17A. The first GC is equipped with a SE52 column and the second one with a modified β -cyclodextrin chiral capillary column. Temperature program: 50 °C (6 min), 50–90 °C at 2 °C/min, 90 °C (20 min); 90–180 °C at 2 °C/min, 180 °C (10 min); injector temperature 250 °C, detector temperature 280 °C, carrier gas: He; injection system: split (ratio 1:150).

NMR spectra were recorded using a Bruker DPX-400 Avance Spectrometer (400 MHz for ^1H , 100 MHz for ^{13}C).

3. Results and discussion

A native strain was isolated from *Eucalyptus* leaves in the presence of **1** as the sole carbon and energy source. This microorganism was identified as *Aspergillus terreus* by our laboratory.

This strain produced four oxygenated derivatives from **1** (see Scheme 2) that were identified as 2-*exo*-hydroxy-1,8-cineole **2**, 2-*endo*-hydroxy-1,8-cineole **3**, 3-*exo*-hydroxy-1,8-cineole **4** and 3-*endo*-hydroxy-1,8-cineole **5**.

The major product (**2**) was identified by comparison with the literature data and co-injection in GC with the synthetic **2**, prepared from alpha-terpineol according to bibliography [24].

The isolated amount for compounds **3** and **5** was not enough to correctly identify these products by NMR, therefore we prepared

Download English Version:

<https://daneshyari.com/en/article/70644>

Download Persian Version:

<https://daneshyari.com/article/70644>

[Daneshyari.com](https://daneshyari.com)