Micro-oxygen bioanode: an efficient strategy for enhancement of phenol degradation and current generation in mix-cultured MFCs

Li-Hui Yang, Ting-Ting Zhu, Wei-Wei Cai, Muhammad Rizwan Haider, Hong-Cheng Wang, Hao-Yi Cheng, Ai-Jie Wang

PII: S0960-8524(18)30922-2
DOI: https://doi.org/10.1016/j.biortech.2018.07.025
Reference: BITE 20155

To appear in: *Bioresource Technology*

Received Date: 23 May 2018
Revised Date: 3 July 2018
Accepted Date: 6 July 2018

Please cite this article as: Yang, L-H., Zhu, T-T., Cai, W-W., Haider, M.R., Wang, H-C., Cheng, H-Y., Wang, A-J., Micro-oxygen bioanode: an efficient strategy for enhancement of phenol degradation and current generation in mix-cultured MFCs, *Bioresource Technology* (2018), doi: https://doi.org/10.1016/j.biortech.2018.07.025

Micro-oxygen bioanode: an efficient strategy for enhancement of phenol degradation and current generation in mix-cultured MFCs

Li-Hui Yang [a,b] , Ting-Ting Zhu [a], Wei-Wei Cai [c], Muhammad Rizwan Haider [a,b], Hong-Cheng Wang [a,b], Hao-Yi Cheng [a], Ai-Jie Wang*[a,c]

[a] *Key Laboratory of Environmental Biotechnology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, P. R. China.*

[b] *University of Chinese Academy of Sciences, Beijing, 100049, P. R. China.*

[c] *State Key Laboratory of Urban Water Resource and Environment, Harbin Institute of Technology, Harbin 150090, PR China*

* Corresponding author: Key Laboratory of Environmental Biotechnology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, P.R. China; E-mail address: ajwang@rcees.ac.cn (Ai-Jie Wang)