Accepted Manuscript

Temperature-phased anaerobic digestion of food waste: A comparison with single-stage digestions based on performance and energy balance

Benyi Xiao, Yu Qin, Wenzhe Zhang, Jing Wu, Hong Qiang, Junxin Liu, Yu-You Li

PII: S0960-8524(17)31913-2

DOI: https://doi.org/10.1016/j.biortech.2017.10.084

Reference: BITE 19125

To appear in: Bioresource Technology

Received Date: 6 September 2017 Revised Date: 23 October 2017 Accepted Date: 24 October 2017



Please cite this article as: Xiao, B., Qin, Y., Zhang, W., Wu, J., Qiang, H., Liu, J., Li, Y-Y., Temperature-phased anaerobic digestion of food waste: A comparison with single-stage digestions based on performance and energy balance, *Bioresource Technology* (2017), doi: https://doi.org/10.1016/j.biortech.2017.10.084

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

1	Temperature-phased anaerobic digestion of food waste: A
2	comparison with single-stage digestions based on performance and
3	energy balance
4	Benyi Xiao ^{1,2} , Yu Qin ³ , Wenzhe Zhang ^{1,2} , Jing Wu ⁴ , Hong Qiang ⁵ , Junxin Liu ^{1,2} ,
5	Yu-You Li ^{3,4*}
6	1. Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing
7	100085, China; 2. University of Chinese Academy of Sciences, Beijing 100049, China; 3.
8	Department of Civil and Environmental Engineering, Graduate School of Engineering, Tohoku
9	University, 6-6-06 Aoba, Aramaki-Aza, Aoba-Ku, Sendai, Miyagi 980-8579, Japan; 4.
10	Department of Frontier Science for Advanced Environment, Graduate School of Environmental
11	Sciences, Tohoku University, 6-6-20 Aoba, Aramaki-Aza, Aoba-Ku, Sendai, Miyagi 980-8579,
12	Japan. 5. College of Resources and Environment, Northwest A&F University, Yangling 712100,
13	China.
14	
15	Abstract
16	
17	The temperature-phased anaerobic digestion (TPAD) of food waste was studied for
18	the purpose of comparing with single-stage mesophilic and thermophilic anaerobic
19	digestion. The biogas and methane yields in the TPAD during the steady period was
	*Corresponding author. Tel.: +81-22-795-7464; Fax: +81-22-795-7465. E-mail address: gyokuyu.ri.a5@tohoku.ac.jp

Download English Version:

https://daneshyari.com/en/article/7068821

Download Persian Version:

https://daneshyari.com/article/7068821

<u>Daneshyari.com</u>