

## Accepted Manuscript

Production of monosaccharides from napier grass by hydrothermal process with phosphoric acid

Eri Takata, Ken Tsutsumi, Yuji Tsutsumi, Kenji Tabata

PII: S0960-8524(13)00902-4  
DOI: <http://dx.doi.org/10.1016/j.biortech.2013.05.112>  
Reference: BITE 11899

To appear in: *Bioresource Technology*

Received Date: 12 March 2013  
Revised Date: 27 May 2013  
Accepted Date: 28 May 2013

Please cite this article as: Takata, E., Tsutsumi, K., Tsutsumi, Y., Tabata, K., Production of monosaccharides from napier grass by hydrothermal process with phosphoric acid, *Bioresource Technology* (2013), doi: <http://dx.doi.org/10.1016/j.biortech.2013.05.112>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



## **Production of monosaccharides from napier grass by hydrothermal process with phosphoric acid**

Eri Takata <sup>1,2</sup>, Ken Tsutsumi <sup>1</sup>, Yuji Tsutsumi <sup>2,\*</sup>, Kenji Tabata <sup>1</sup>.

<sup>1</sup>*Department of Applied Chemistry, Faculty of Engineering, University of Miyazaki, 1-1, Gakuen-kibanadai-nishi, Miyazaki, 889-2192, Japan*

<sup>2</sup>*Department of agro-environmental Sciences, Faculty of Agriculture, Kyushu University, 6-10-1, Hakozaki Higashi-ku, Fukuoka, 812-8581, Japan*

\* Corresponding author. Tel: +81-92-642-2988; Fax: +81-92-642-2988; Email address: [y-tsutsu@agr.kyushu-u.ac.jp](mailto:y-tsutsu@agr.kyushu-u.ac.jp)

Download English Version:

<https://daneshyari.com/en/article/7081708>

Download Persian Version:

<https://daneshyari.com/article/7081708>

[Daneshyari.com](https://daneshyari.com)