

Contents lists available at ScienceDirect



Journal of King Saud University – Science

journal homepage: www.sciencedirect.com

Existence and uniqueness results for a fractional Riemann–Liouville nonlocal thermistor problem on arbitrary time scales

Moulay Rchid Sidi Ammi^a, Delfim F.M. Torres^{b,*}

^a AMNEA Group, Department of Mathematics, Faculty of Sciences and Techniques, Moulay Ismail University, B.P. 509, Errachidia, Morocco

^b Center for Research & Development in Mathematics and Applications (CIDMA), Department of Mathematics, University of Aveiro, 3810-193 Aveiro, Portugal

ARTICLE INFO

Article history:

Received 16 December 2016

Accepted 14 March 2017

Available online xxxxx

MSC:

26A33

26E70

35B09

45M20

Keywords:

Fractional Riemann–Liouville derivatives
 Nonlocal thermistor problem on time scales
 Fixed point theorem
 Dynamic equations
 Positive solutions

ABSTRACT

Using a fixed point theorem in a proper Banach space, we prove existence and uniqueness results of positive solutions for a fractional Riemann–Liouville nonlocal thermistor problem on arbitrary nonempty closed subsets of the real numbers.

© 2017 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The calculus on time scales is a recent area of research introduced by [Aulbach and Hilger \(1990\)](#), unifying and extending the theories of difference and differential equations into a single theory. A time scale is a model of time, and the theory has found important applications in several contexts that require simultaneous modeling of discrete and continuous data. It is under strong current research in areas as diverse as the calculus of variations, optimal control, economics, biology, quantum calculus, communication networks and robotic control. The interested reader is referred to [Agarwal and Bohner \(1999\)](#), [Agarwal et al. \(2002\)](#), [Bohner](#)

and [Peterson \(2001a,b\)](#), [Martins and Torres \(2009\)](#), [Ortigueira et al. \(2016\)](#) and references therein.

On the other hand, many phenomena in engineering, physics and other sciences, can be successfully modeled by using mathematical tools inspired by the fractional calculus, that is, the theory of derivatives and integrals of noninteger order. See, for example, [Gaul et al. \(1991\)](#), [Hilfer \(2000\)](#), [Kilbas et al. \(2006\)](#), [Sabatier et al. \(2007\)](#), [Samko et al. \(1993\)](#) and [Srivastava and Saxena \(2001\)](#). This allows one to describe physical phenomena more accurately. In this line of thought, fractional differential equations have emerged in recent years as an interdisciplinary area of research ([Abbas et al., 2012](#)). The nonlocal nature of fractional derivatives can be utilized to simulate accurately diversified natural phenomena containing long memory ([Debbouche and Torres, 2015](#); [Machado et al., 2011](#)).

A thermistor is a thermally sensitive resistor whose electrical conductivity changes drastically by orders of magnitude, as the temperature reaches a certain threshold. Thermistors are used as temperature control elements in a wide variety of military and industrial equipment, ranging from space vehicles to air conditioning controllers. They are also used in the medical field, for localized and general body temperature measurement; in meteorology, for

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

* Corresponding author.

E-mail addresses: rachidsidiammi@yahoo.fr (M.R. Sidi Ammi), delfim@ua.pt (D.F. M. Torres).

<http://dx.doi.org/10.1016/j.jksus.2017.03.004>

1018–3647/© 2017 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article in press as: Sidi Ammi, M.R., Torres, D.F.M.. Journal of King Saud University – Science (2017), <http://dx.doi.org/10.1016/j.jksus.2017.03.004>

weather forecasting; as well as in chemical industries, as process temperature sensors (Kwok, 1995; Maclen, 1979).

Throughout the reminder of the paper, we denote by \mathbb{T} a time scale, which is a nonempty closed subset of \mathbb{R} with its inherited topology. For convenience, we make the blanket assumption that t_0 and T are points in \mathbb{T} . Our main concern is to prove existence and uniqueness of solution to a fractional order nonlocal thermistor problem of the form

$$\begin{aligned} {}_{t_0}^{\mathbb{T}}D_t^{2\alpha}u(t) &= \frac{\lambda f(u)}{\left(\int_{t_0}^T f(u) \Delta x\right)^2}, \quad t \in (t_0, T), \\ {}_{t_0}^{\mathbb{T}}I_t^\beta u(t_0) &= 0, \quad \forall \beta \in (0, 1), \end{aligned} \tag{1}$$

under suitable conditions on f as described below. We assume that $\alpha \in (0, 1)$ is a parameter describing the order of the fractional derivative; ${}_{t_0}^{\mathbb{T}}D_t^{2\alpha}$ is the left Riemann–Liouville fractional derivative operator of order 2α on ${}_{t_0}^{\mathbb{T}}I_t^\beta$ is the left Riemann–Liouville fractional integral operator of order β defined on \mathbb{T} by Benkhettou et al. (2016b). By u , we denote the temperature inside the conductor; $f(u)$ is the electrical conductivity of the material.

In the literature, many existence results for dynamic equations on time scales are available (Dogan, 2013a; Dogan, 2013b). In recent years, there has been also significant interest in the use of fractional differential equations in mathematical modeling (Aghababa, 2015; Ma et al., 2016; Yu et al., 2016). However, much of the work published to date has been concerned separately, either by the time-scale community or by the fractional one. Results on fractional dynamic equations on time scales are scarce (Ahmadkhanlu and Jahanshahi, 2012).

In contrast with our previous works, which make use of fixed point theorems like the Krasnoselskii fixed point theorem, the fixed point index theory, and the Leggett–Williams fixed point theorem, to obtain several results of existence of positive solutions to linear and nonlinear dynamic equations on time scales, and recently also to fractional differential equations (Sidi Ammi et al., 2012; Sidi Ammi and Torres, 2012b, 2013; Souahi et al., 2016); here we prove new existence and uniqueness results for the fractional order nonlocal thermistor problem on time scales (1), putting together time scale and fractional domains. This seems to be quite appropriate from the point of view of practical applications (Machado et al., 2015; Nwaeze and Torres, 2017; Ortigueira et al., 2016).

The rest of the article is arranged as follows. In Section 2, we state preliminary definitions, notations, propositions and properties of the fractional operators on time scales needed in the sequel. Our main aim is to prove existence of solutions for (1) using a fixed point theorem and, consequently, uniqueness. This is done in Section 3: see Theorems 3.2 and 3.6.

2. Preliminaries

In this section, we recall fundamental definitions, hypotheses and preliminary facts that are used through the paper. For more details, see the seminal paper Benkhettou et al. (2016b). From physical considerations, we assume that the electrical conductivity is bounded (Antontsev and Chipot, 1994). Precisely, we consider the following assumption:

(H1). Function $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ of problem (1) is Lipschitz continuous with Lipschitz constant L_f such that $c_1 \leq f(u) \leq c_2$, with c_1 and c_2 two positive constants.

We deal with the notions of left Riemann–Liouville fractional integral and derivative on time scales, as proposed in Benkhettou et al. (2016b), the so called BHT fractional calculus on time scales

(Nwaeze and Torres, 2017). The corresponding right operators are obtained by changing the limits of integrals from a to t into t to b . For local approaches to fractional calculus on arbitrary time scales we refer the reader to Benkhettou et al. (2015, 2016a). Here we are interested in nonlocal operators, which are the ones who make sense with respect to the thermistor problem (Sidi Ammi and Torres, 2008, 2012a). Although we restrict ourselves to the delta approach on time scales, similar results are trivially obtained for the nabla fractional case (Girejko and Torres, 2012).

Definition 2.1 (Riemann–Liouville fractional integral on time scales (Benkhettou et al., 2016b)). Let \mathbb{T} be a time scale and $[a, b]$ an interval of \mathbb{T} . Then the left fractional integral on time scales of order $0 < \alpha < 1$ of a function $g : \mathbb{T} \rightarrow \mathbb{R}$ is defined by

$${}_{a}^{\mathbb{T}}I_t^\alpha g(t) = \int_a^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} g(s) \Delta s,$$

where Γ is the Euler gamma function.

The left Riemann–Liouville fractional derivative operator of order α on time scales is then defined using Definition 2.1 of fractional integral.

Definition 2.2 (Riemann–Liouville fractional derivative on time scales (Benkhettou et al., 2016b)). Let \mathbb{T} be a time scale, $[a, b]$ an interval of \mathbb{T} , and $0 < \alpha < 1$. Then the left Riemann–Liouville fractional derivative on time scales of order α of a function $g : \mathbb{T} \rightarrow \mathbb{R}$ is defined by

$${}_{a}^{\mathbb{T}}D_t^\alpha g(t) = \left(\int_a^t \frac{(t-s)^{-\alpha}}{\Gamma(1-\alpha)} g(s) \Delta s \right)^\Delta.$$

Remark 2.3. If $\mathbb{T} = \mathbb{R}$, then we obtain from Definitions 2.1 and 2.2, respectively, the usual left Riemann–Liouville fractional integral and derivative.

Proposition 2.4 (See Benkhettou et al. (2016b)). Let \mathbb{T} be a time scale, $g : \mathbb{T} \rightarrow \mathbb{R}$ and $0 < \alpha < 1$. Then

$${}_{a}^{\mathbb{T}}D_t^\alpha g = \Delta \circ {}_{a}^{\mathbb{T}}I_t^{1-\alpha} g.$$

Proposition 2.5 (See Benkhettou et al. (2016b)). If $\alpha > 0$ and $g \in C([a, b])$, then

$${}_{a}^{\mathbb{T}}D_t^\alpha \circ {}_{a}^{\mathbb{T}}I_t^\alpha g = g.$$

Proposition 2.6 (See Benkhettou et al. (2016b)). Let $g \in C([a, b])$ and $0 < \alpha < 1$. If ${}_{a}^{\mathbb{T}}I_t^{1-\alpha} u(a) = 0$, then

$${}_{a}^{\mathbb{T}}I_t^\alpha \circ {}_{a}^{\mathbb{T}}D_t^\alpha g = g.$$

Theorem 2.7 (See Benkhettou et al. (2016b)). Let $g \in C([a, b])$, $\alpha > 0$, and ${}_{a}^{\mathbb{T}}I_t^\alpha([a, b])$ be the space of functions that can be represented by the Riemann–Liouville Δ -integral of order α of some $C([a, b])$ -function. Then,

$$g \in {}_{a}^{\mathbb{T}}I_t^\alpha([a, b])$$

if and only if

$${}_{a}^{\mathbb{T}}I_t^{1-\alpha} g \in C^1([a, b])$$

and

$${}_{a}^{\mathbb{T}}I_t^{1-\alpha} g(a) = 0.$$

Download English Version:

<https://daneshyari.com/en/article/7216582>

Download Persian Version:

<https://daneshyari.com/article/7216582>

[Daneshyari.com](https://daneshyari.com)