



## Short Communication

## Effect of the *Lippia alba* (Mill.) N.E. Brown essential oil and its main constituents, citral and limonene, on the tracheal smooth muscle of rats



Poliana M.M. Carvalho<sup>a</sup>, Cícero A.F. Macêdo<sup>a</sup>, Tiago F. Ribeiro<sup>a</sup>, Andressa A. Silva<sup>a</sup>,  
Renata E.R. Da Silva<sup>a</sup>, Luís P. de Moraes<sup>a</sup>, Marta R. Kerntopf<sup>b</sup>, Irwin R.A. Menezes<sup>c</sup>,  
Roseli Barbosa<sup>a,\*</sup>

<sup>a</sup> Physiopharmacology of Excitability Cell Laboratory, Department of Chemical Biology Regional University of Cariri, Campus of Pimenta, 63105-010, Crato, CE, Brazil

<sup>b</sup> Pharmacology of Natural Products Laboratory, Regional University of Cariri-Campus of Pimenta, 63105-010, Crato, CE, Brazil

<sup>c</sup> Pharmacology and Molecular Chemistry Laboratory, Department of Chemical Biology, Regional University of Cariri-Campus of Pimenta, 63105-010, Crato, CE, Brazil

## ARTICLE INFO

## Key-words:

*Lippia alba*  
Essential oil  
Citral  
Smooth muscle  
Limonene

## ABSTRACT

The *Lippia alba* (Mill.) N.E. Brown (Verbenaceae) species, has effects sedative, analgesic and spasmolytic properties. This study had as its main objective to evaluate the essential oil of *L. alba* (EOLa) effect and that of its main constituents, citral and limonene, over tracheal smooth muscle from Wistar rats. EOLa, citral and limonene promoted relaxation of tracheal smooth muscle in contractions induced by potassium ( $60 \text{ mM K}^+$ ), presenting an  $EC_{50}$  of  $148 \pm 7 \mu\text{g/mL}$  for the EOLa,  $136 \pm 7 \mu\text{g/mL}$  for citral and  $581 \pm 7 \mu\text{g/mL}$  for limonene. In contractions induced by Acetylcholine (Ach;  $10 \mu\text{M}$ ) the  $EC_{50}$  for the EOLa and citral were of  $731 \pm 5 \mu\text{g/mL}$  and  $795 \pm 9 \mu\text{g/mL}$ , respectively. In preparations pre-incubated with  $1000 \mu\text{g/mL}$  of the EOLa and citral, both agents were found to block the influx of  $\text{BaCl}_2$  by VOCCs. This study demonstrated that the EOLa and its main component citral present antispasmodic effect over tracheal smooth muscle of rats.

## 1. Introduction

The species *Lippia alba* (Mill.) N. E. Brown (Verbenaceae) is an aromatic shrub native to South America, Central America and Africa. In Brasil, *L. alba* is known popularly as “lemon balm”, “false melissa” and “field rosemary” and it is readily used in the form of infusion or decoction of the leaves for the treatment of cold, bronchitis, cough, asthma and intestinal disorders [1]. Pharmacological activities have also been verified for the species: antioxidant, sedative [2], anxiolytic [3] and antispasmodic [4].

Citral and limonene are the main components of the essential oil of *L. alba*. Citral is a natural mixture of aldeidos, being the geranial ((E)-3,7-dimethyl-2,6-octadienal) and the Neral ((Z)-3,7-dimethyl-2,6-octadienal), where the chemotype under study is classified as chemotype II (MATO et al., 1996) [5]. Pharmacological studies attribute to citral the following activities: acaricide, insecticide and anxiolytic, [6] as well as sedative [7]. Limonene, (R)4-isoprenyl-1-methyl-cyclohexene, is a monocyclic monoterpene present in the structure of many plants such as *Mentha* spp. and citrus plants. Pharmacologically, activities such as: antifungal, antibacterial, antitumor, acaricide, insecticide and repellent activities, have been attributed to limonene [8–12].

Currently, the pharmaceutical industry have been searching for

substances in natural products which can improve the treatment of many diseases, among which, asthma, allergic rhinitis and chronic obstructive pulmonary disease, which affect a large part of the world population, stand out. Due to the lack of studies with *L. alba* on the upper respiratory tract muscles, the present study sought to evaluate the effect of the essential oil of *Lippia alba* (EOLa) and its main constituents, citral and limonene, on the tracheal smooth muscle contractions of Wistar rats.

## 2. Materials and methods

## 2.1. Botanic material

The botanical material was obtained at the UFC experimental farm by Dr. Sergio Horta (experimental farm of the Federal University of Ceará, Brazil). The essential oil extracted from a *Lippia alba* (Mill.) N. E. Brown sample was chemically analyzed in the Laboratory of Natural Products and the Technological Development Park (PADETEC) of the Federal University of Ceará, Brazil, with the following compounds being detected: citral 75.92% [Geraniol (41.81%) and neral (34.11%)], limonene (9.85%), carvone (8.92%), gamma-terpinene (2.05%), cymene (1.02%). As observed in work by Sousa et al. [13].

\* Corresponding author at: Physiopharmacology of Excitability Cell Laboratory, University Regional of Cariri, Campus of Pimenta, 63105-010, Crato, CE, Brazil.  
E-mail address: [roseli.barbosa@urca.br](mailto:roseli.barbosa@urca.br) (R. Barbosa).

## 2.2. Substances and solutions

The nutrient solution containing the following composition in mM: NaCl = 136; KCl = 5.0; MgCl<sub>2</sub> = 0.98; NaH<sub>2</sub>PO<sub>4</sub> = 0.36; NaHCO<sub>3</sub> = 11.9; CaCl<sub>2</sub> = 2.0 and C<sub>6</sub>H<sub>12</sub>O<sub>6</sub> = 5.5, was maintained at 37 °C and the pH was adjusted to 7.4, remaining in stabilization for 1 h. The EOLA, citral and limonene, were prepared as a solution, diluted directly into Tyrode and Tween (3%). The calcium-free solution or “zero calcium” (0 Ca<sup>2+</sup>) was produced without CaCl<sub>2</sub> in the Tyrode solution and an addition of 0.2 mM EGTA. All salts and reagents used were of analytical grade and purity obtained from the company Sigma-Aldrich (St. Louis, Missouri, USA). To confirm whether smooth muscle relaxation is dependent on voltage-operated cation channels (VOCCs), BaCl<sub>2</sub> and nifedipine (1 μM) were used.

## 2.3. Animals and experimental procedures

Male Wistar rats (*Rattus norvegicus*) with a body mass of 200–300 g, obtained from the Central Biotechnology of the Regional University of Cariri-URCA, Brazil, were used. The animals were kept under constant humidity and temperature conditions of 23 ± 2 °C, in a twelve hours light/dark cycle, with access to water and ration *ad libitum*, and were treated according to the Brazilian College of Animal Experimentation (COBEA), Brazil. The study was approved by the Committee on Ethics in the Use of Animals (CEUA), registered under the protocol number: under No. 24/2012.2/2012.

The animals were euthanised in a CO<sub>2</sub> chamber, followed by the dissection of the trachea, which was sectioned into circular transverse segments of approximately 3 to 4 mm in length, which were then mounted in isolated organ bath tubs with a capacity for 10 mL of modified Tyrode nutrient solution, maintained under continuous aeration by O<sub>2</sub> bubbling, at 37 °C and pH 7.4 for 60 min. Consequently, the contracting agonists KCl (60 mM) and ACh (10 μM) were added to the organ baths in distinct experiments and followed by a crescent and cumulative addition of the EOLA, citral and limonene separately.

The data were presented as the mean ± S.E.M. and N, where N represents the number of experiments and S.E.M. means the standard error of the mean. The software Sigma Plot 11.0 was used for statistical analysis and the production of graphs. The results considered statistically significant had a null hypothesis probability of less than 5% (p < 0.05). The Analysis of Variance test (*one-way ANOVA*), followed by the *Holm-Sidak* multiple comparisons test when appropriate, were used. For the calculation of the EC<sub>50</sub>, logarithmic interpolation was performed, which was considered as the concentration of the substance which is able to produce 50% of its inhibition or its maximum effect, and the calculations were performed for each experiment.

## 3. Results

In the assessment of basal tonus tracheal, the increasing and cumulative concentrations of EOLA, citral and limonene (1–1000 μg/mL), were added to the tracheal rings of Wistar rats at baseline. The results showed that there was no statistically significant relaxant or contracting effect on the basal tonus of the tracheal ring preparations (p > 0.05, *one-way ANOVA*). In preparations pre-contracted with the KCl (60 mM), increasing concentrations of the EOLA, citral and limonene (1–1000 μg/mL), promoted concentration-dependent relaxation, where significant effects were observed in the concentrations ≥ 30 μg/mL for the EOLA, ≥ 30 μg/mL for citral and ≥ 600 μg/mL for limonene, presenting with an EC<sub>50</sub> of 148 ± 7 μg/mL for the EOLA, 136 ± 7 μg/mL for citral and 581 ± 7 μg/mL for limonene, with these being statistically significant (Fig. 1) (p < 0.001, *one-way ANOVA* followed by *Holm-Sidak*).

When investigating the activity of the EOLA, citral and limonene (1–3000 μg/mL), on the contractions evoked by ACh (10 μM), there was a myorelaxant activity for EOLA and citral, and its effects were




Fig. 1. Concentration-effect curve of OELA, citral and limonene (1–1000 μg/mL), under contraction evoked by KCl (60 mM) in trachea isolated from rats. Values are expressed as mean ± SEM, (p < 0.05, *one-way ANOVA* followed by *Holm-Sidak*), where N represents the number of experiments.




Fig. 2. Concentration-effect curve of OELA, citral and limonene (1–3000 μg/mL), in contraction evoked by ACh (10 μM) in trachea isolated from rats. Values are expressed as mean ± S.E.M. (p < 0.05, *one-way ANOVA* followed by *Holm-Sidak*), where N represents the number of experiments.

significant in concentrations ≥ 300 μg/mL for the EOLA and ≥ 600 μg/mL for citral, presenting with an EC<sub>50</sub> of 731 ± 5 μg/mL and 795 ± 9 μg/mL for the EOLA and citral, respectively (Fig. 2) (p < 0.001, *one-way ANOVA*, followed by *Holm-sidak*). Limonene was not able to produce a statistically significant myorelaxant effect (p > 0.05, *one-way ANOVA*).

According to the results, we can observe that the EOLA and citral have their most pronounced action in the electromechanical pathway, when comparing its results with the pharmacological route, thus demonstrating that the EOLA and citral act predominantly on VOCCs channels.

To investigate whether this smooth muscle relaxation occurs through L-type calcium channels, we used cumulative BaCl<sub>2</sub> (1–30 mM) concentrations, which is an ion with selectivity for VOCCs [14].

In preparations pre-incubated with 1000 μg/mL of the EOLA and citral, it was possible to verify that both agents blocked the influx of BaCl<sub>2</sub> by the VOCCs, promoting smooth muscle relaxation, with a similar behavior to that of nifedipine (1 μM), this being a voltage-dependent calcium channel blocker. Whereas in preparations induced with 1000 μM limonene, the contraction induced by BaCl<sub>2</sub> was allowed, demonstrating that limonene does not block L-type calcium channels (Fig. 3).

Download English Version:

<https://daneshyari.com/en/article/7234938>

Download Persian Version:

<https://daneshyari.com/article/7234938>

[Daneshyari.com](https://daneshyari.com)