

Accepted Manuscript

Title: Resting-state connectivity of the amygdala predicts response to cognitive behavioral therapy in obsessive compulsive disorder

Author: Martin Göttlich Ulrike M. Krämer Andreas Kordon
Fritz Hohagen Bartosz Zurowski

PII: S0301-0511(15)30053-3
DOI: <http://dx.doi.org/doi:10.1016/j.biopsycho.2015.09.004>
Reference: BIOPSY 7091

To appear in:

Received date: 22-7-2014
Revised date: 31-8-2015
Accepted date: 14-9-2015

Please cite this article as: Göttlich, Martin, Krämer, Ulrike M., Kordon, Andreas, Hohagen, Fritz, Zurowski, Bartosz, Resting-state connectivity of the amygdala predicts response to cognitive behavioral therapy in obsessive compulsive disorder. *Biological Psychology* <http://dx.doi.org/10.1016/j.biopsycho.2015.09.004>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Resting-state connectivity of the amygdala predicts response to cognitive behavioral therapy in obsessive compulsive disorder

Abbreviated title: Connectivity and therapy outcome in OCD

Martin Göttlich¹, Ulrike M. Krämer¹, Andreas Kordon², Fritz Hohagen², Bartosz Zurowski^{2,3}

¹Dept. of Neurology, University of Lübeck, Lübeck, Germany

²Dept. of Psychiatry, University of Lübeck, Lübeck, Germany

³University of Hamburg, Institute for Systems Neuroscience, Germany

Correspondence and requests for materials should be addressed to
Martin Göttlich (martin.goettlich@neuro.uni-luebeck.de)
Universitätsklinikum Schleswig-Holstein - Campus Lübeck
Klinik für Neurologie
Ratzeburger Allee 160
23562 Lübeck
Tel.: +49 451 317931313

Highlights

- We investigated voxel-wise whole-brain resting-state connectivity in OCD patients.
- Functional data was acquired prior to an 8-week cognitive behavioral therapy (CBT).
- Connectivity of the basolateral amygdala predicted the outcome of CBT.

Download English Version:

<https://daneshyari.com/en/article/7278554>

Download Persian Version:

<https://daneshyari.com/article/7278554>

[Daneshyari.com](https://daneshyari.com)