

## Accepted Manuscript

Angiotensin receptor blocker irbesartan reduces stress-induced intestinal inflammation via AT1a signaling and ACE2-dependent mechanism in mice

Maimaiti Yisireyili, Yasuhiro Uchida, Koji Yamamoto, Takayuki Nakayama, Xian Wu Cheng, Tadashi Matsushita, Shigeo Nakamura, Toyooki Murohara, Kyosuke Takeshita

PII: S0889-1591(17)30511-1  
DOI: <https://doi.org/10.1016/j.bbi.2017.11.010>  
Reference: YBRBI 3281

To appear in: *Brain, Behavior, and Immunity*

Received Date: 7 June 2017  
Revised Date: 2 November 2017  
Accepted Date: 15 November 2017

Please cite this article as: Yisireyili, M., Uchida, Y., Yamamoto, K., Nakayama, T., Cheng, X.W., Matsushita, T., Nakamura, S., Murohara, T., Takeshita, K., Angiotensin receptor blocker irbesartan reduces stress-induced intestinal inflammation via AT1a signaling and ACE2-dependent mechanism in mice, *Brain, Behavior, and Immunity* (2017), doi: <https://doi.org/10.1016/j.bbi.2017.11.010>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**Angiotensin receptor blocker irbesartan reduces stress-induced intestinal inflammation via AT1a signaling and ACE2-dependent mechanism in mice**

Maimaiti Yisireyili<sup>1,7</sup>, Yasuhiro Uchida<sup>1</sup>, Koji Yamamoto<sup>2</sup>, Takayuki Nakayama<sup>3</sup>,  
5 Xian Wu Cheng<sup>1</sup>, Tadashi Matsushita<sup>4,5</sup>, Shigeo Nakamura<sup>6</sup>, Toyoaki Murohara<sup>1</sup>, and  
Kyosuke Takeshita<sup>1,4</sup>

<sup>1</sup>Department of Cardiology, Nagoya University Graduate School of Medicine, Nagoya, Japan. Departments of <sup>4</sup>Clinical Laboratory, <sup>5</sup>Blood Transfusion, and <sup>6</sup>Pathology,  
10 Nagoya University Hospital, Nagoya, Japan. <sup>2</sup>Department of Transfusion Medicine and Cell Therapy, Saitama Medical Centre, Saitama Medical University, Kawagoe, Japan. <sup>3</sup>Department of Blood Transfusion, Aichi Medical University Hospital, Nagakute, Japan. <sup>7</sup>Department of Minimally Invasive Hernia and Abdominal Surgery, People's Hospital of Xinjiang Uygur Autonomous Region, Urumqi 830000, Xinjiang  
15 Uygur Autonomous Region, China.

*Running title:* ARB reduces stress-induced intestinal inflammation.

*Abstract word count:* 253

*Total word count:* 8129

**Correspondence to:**

A/Prof. Kyosuke Takeshita, MD, PhD, FAHA.  
Department of Cardiology  
Nagoya University Graduate School of Medicine  
65 Tsurumai-cho Nagoya Aichi 466-8550, Japan.

Download English Version:

<https://daneshyari.com/en/article/7279394>

Download Persian Version:

<https://daneshyari.com/article/7279394>

[Daneshyari.com](https://daneshyari.com)