
Accepted Manuscript

Induction of suppressor of cytokine signaling 3 via HSF-1-HSP70-TLR4 axis attenuates neuroinflammation and ameliorates postoperative pain

Yi-Xin Fan, Cheng Qian, Bingqian Liu, Chaoyu Wang, Haijiao Liu, Xiuxiu Pan, Peng Teng, Liang Hu, Guangqin Zhang, Yuan Han, Mi Yang, Xue-Feng Wu, Wen-Tao Liu

PII: DOI: Reference:	S0889-1591(17)30459-2 https://doi.org/10.1016/j.bbi.2017.10.006 YBRBI 3250
To appear in:	Brain, Behavior, and Immunity
Received Date:	1 April 2017

Received Date:1 April 2017Revised Date:30 September 2017Accepted Date:6 October 2017



Please cite this article as: Fan, Y-X., Qian, C., Liu, B., Wang, C., Liu, H., Pan, X., Teng, P., Hu, L., Zhang, G., Han, Y., Yang, M., Wu, X-F., Liu, W-T., Induction of suppressor of cytokine signaling 3 via HSF-1-HSP70-TLR4 axis attenuates neuroinflammation and ameliorates postoperative pain, *Brain, Behavior, and Immunity* (2017), doi: https://doi.org/10.1016/j.bbi.2017.10.006

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Induction of suppressor of cytokine signaling 3 via HSF-1-HSP70-TLR4 axis attenuates neuroinflammation and ameliorates postoperative pain

Yi-Xin Fan^{a, 1}, Cheng Qian^{a, 1}, Bingqian Liu^c, Chaoyu Wang^b, Haijiao Liu^d, Xiuxiu Pan^b, Peng Teng^b, Liang Hu^b, Guangqin Zhang^d, Yuan Han^e, Mi Yang^f, Xue-Feng Wu^g, and Wen-Tao Liu^{a, b *}

^a Department of Pharmacy, Sir Run Run Shaw Hospital Affiliated to Nanjing Medical University, Jiangsu, 211166, People's Republic of China

^b Jiangsu Key Laboratory of Neurodegeneration, Department of Pharmacology, Nanjing Medical University, Nanjing, China

^c Department of ophthalmology, The First Affiliated Hospital with Nanjing Medical University, Jiangsu, China

^d Research Division of Pharmacology, China Pharmaceutical University, Nanjing, China

^e Jiangsu Province Key Laboratory of Anesthesiology, Xuzhou Medical College, Xuzhou, Jiangsu, China

^f The Comprehensive Cancer Center of Drum Tower Hospital, Medical School of Nanjing University & Clinical Cancer Institute of Nanjing University, Nanjing, China ^g State Key Laboratory of Pharmaceutical Biotechnology and Collaborative Innovation Center of Chemistry for Life Sciences, School of Life Sciences, Nanjing University, Jiangsu, China

*Correspondence authors at: Department of Pharmacy, Sir Run Run Shaw Hospital Affiliated to Nanjing Medical University, Jiangsu, 211166, People's Republic of China.

Jiangsu Key Laboratory of Neurodegeneration, Department of Pharmacology, Nanjing Medical University, Nanjing, China.

E-mail addresses: painresearch@njmu.edu.cn.

¹These authors contributed equally to this work.

Download English Version:

https://daneshyari.com/en/article/7279606

Download Persian Version:

https://daneshyari.com/article/7279606

Daneshyari.com