# Accepted Manuscript


Title: Sensory aspects of Tourette syndrome

Authors: Joanna H. Cox, Stefano Seri, Andrea E. Cavanna

PII: S0149-7634(18)30079-4
DOI: https://doi.org/10.1016/j.neubiorev.2018.03.016
Reference: NBR 3074

To appear in:

Received date: 2-2-2018
Revised date: 14-3-2018
Accepted date: 15-3-2018

Please cite this article as: Cox JH, Seri S, Cavanna AE, Sensory aspects of Tourette syndrome, *Neuroscience and Biobehavioral Reviews* (2010), https://doi.org/10.1016/j.neubiorev.2018.03.016

*Invited Article*

# Sensory Aspects of Tourette Syndrome

Joanna H Cox[1], Stefano Seri[2,3], Andrea E Cavanna[2,4,5*]

[1] University Hospitals Birmingham NHS Foundation Trust, Birmingham, United Kingdom
[2] School of Life and Health Sciences, Aston Brain Centre, Aston University, Birmingham, United Kingdom
[3] Department of Clinical Neurophysiology, The Birmingham Women's and Children's Hospital NHS Foundation Trust, Birmingham, United Kingdom
[4] Department of Neuropsychiatry, BSMHFT and University of Birmingham, Birmingham, United Kingdom
[5] Sobell Department of Motor Neuroscience and Movement Disorders, Institute of Neurology and University College London, London, United Kingdom

*Correspondence:

Prof Andrea E. Cavanna, MD PhD FRCP FANPA
Department of Neuropsychiatry
The National Centre for Mental Health
25 Vincent Drive
Birmingham B15 2FG
United Kingdom
Email: a.e.cavanna@bham.ac.uk

## Highlights

- Patients with Tourette syndrome (TS) report that their tics have sensory correlates.
- Premonitory urges and multimodal hypersensitivity are common sensory features of TS.
- Both the insula and sensorimotor areas might be involved in the pathophysiology of TS.