



## Research report

# The salt content of products from popular fast-food chains in Costa Rica <sup>☆</sup>



Katrina Heredia-Blonval <sup>a,\*</sup>, Adriana Blanco-Metzler <sup>a</sup>, Marielos Montero-Campos <sup>a</sup>, Elizabeth K. Dunford <sup>b</sup>

<sup>a</sup> Unit of Nutrition and Health, Instituto Costarricense de Investigación y Enseñanza en Nutrición y Salud (INCIENSA), PO Box 4-2250, Tres Ríos, Costa Rica

<sup>b</sup> The George Institute for Global Health, University of Sydney, PO Box M201, Missenden Rd, Camperdown, NSW 2050, Australia

## ARTICLE INFO

## Article history:

Received 10 October 2013

Received in revised form 31 July 2014

Accepted 20 August 2014

Available online 27 August 2014

## Keywords:

Fast foods

Salt

Sodium

Labeling

## ABSTRACT

Salt is a major determinant of population blood pressure levels. Salt intake in Costa Rica is above levels required for good health. With an increasing number of Costa Ricans visiting fast food restaurants, it is likely that fast-food is contributing to daily salt intake. Salt content data from seven popular fast food chains in Costa Rica were collected in January 2013. Products were classified into 10 categories. Mean salt content was compared between chains and categories. Statistical analysis was performed using Welch ANOVA and Tukey–Kramer HSD tests. Significant differences were found between companies; Subway products had lowest mean salt content (0.97 g/100 g;  $p < 0.05$ ) while Popeye's and KFC had the highest (1.57 g/100 g;  $p < 0.05$ ). Significant variations in mean salt content were observed between categories. Salads had a mean salt content of 0.45 g/100 g while sauces had 2.16 g/100 g ( $p < 0.05$ ). Wide variation in salt content was also seen within food categories. Salt content in sandwiches ranged from 0.5 to 2.1 g/100 g. The high levels and wide variation in salt content of fast food products in Costa Rica suggest that salt reduction is likely to be technically feasible in many cases. With an increasing number of consumers purchasing fast foods, even small improvements in salt levels could produce important health gains.

© 2014 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

## Introduction

Many countries around the world are experiencing a nutrition transition, and Costa Rica is no exception. Globally, populations are changing from a diet based on fresh grains, tubers, fruits and vegetables to a diet based on pre-packaged foods which can often be more energy-dense and higher in total fat, saturated fat, cholesterol, sugar, and sodium (Borbón, Robles, & Huesca, 2010; Fernández, Granados, Sandoval, & Acuña, 2010; Johnson et al., 2010; Marchioni, Claro, Levy, & Monteiro, 2011; Monteiro, Levy, Moreira, Rugani de Castro, & Cannon, 2011; Nielsen & Jopkin, 2003; Paeratakul, Ferdinand, Champagne, Ryan, & Bray, 2003). The globalization of food systems has transformed the traditional diet in developing countries like Costa Rica, due to the increasing availability of processed and fast food (Creel, Sharkey, McIntosh, Anding, & Huber, 2008). This situation has contributed to the spread of obesity and non-communicable diseases (hypertension, diabetes, and cardiovascular problems) (Monge, 2001; O'Donnell, Hoerr, Mendoza, & Tsuei Goh,

2008; Rosenheck, 2008; Schmidt et al., 2005), especially among people of lower socioeconomic status, that are also prone to under-nutrition (Kearney, 2010).

It is well documented that a high consumption of both energy and dietary sodium are associated with the development of high blood pressure (HBP) or hypertension, a leading risk factor for cardiovascular disease (CVD) (Campbell, Jilliian, & Campbell, 2012; Prentice & Jebb, 2003; World Health Organization, 2007). In Costa Rica CVD is the leading cause of death (33%) in both men and women over 30 years of age. One in three adults in Costa Rica suffers from HBP, a number that has risen more than 10% in the last 15 years (Ministerio de Salud, 2011).

The consumption of sodium from all sources in Costa Rica is higher than the World Health Organization recommendation of 5 g per day (World Health Organization, 2012). It is estimated that the contribution of sodium from table salt is 2.8 g per person per day (equivalent to 7 g of sodium chloride or table salt), and although the exact contribution of sodium from processed and fast foods is not known, it is hypothesized to be higher than what is recommended (Ministerio de Salud, 2011). It is also well documented that in Costa Rica families, teens and young adults are the groups most likely to frequent fast food restaurants, (Euromonitor Internacional, 2011) particularly in an effort to save time and increasingly as a dining option with their families and friends (Monge, 2001; Monge, Aragón, Chinnock, Campos, & Colón, 2013; Monge, Smith, Colón,

<sup>☆</sup> Acknowledgements: This research was supported by Canada's International Development Research Centre (Project #106 888), The George Institute for Global Health.

\* Corresponding author.

E-mail address: [katrihe@gmail.com](mailto:katrihe@gmail.com), [kheredia@inciensa.sa.cr](mailto:kheredia@inciensa.sa.cr) (K. Heredia-Blonval).

Aragón, & Herrera, 2013). Most of the fast food chains in Costa Rica were initially established in the USA, highlighting the influence of American culture in the eating habits of Costa Ricans (Bermúdez & Tucker, 2003; Kearney, 2010).

A number of transnational companies around the world have committed to reformulating their products toward healthier formulations and creating new ones with better nutritional quality (less sodium, sugar and fat) (Dunford, 2012). Many transnational fast food chains now operate in Costa Rica, however, studies show that salt in fast foods varies substantially between chains and from one country to another (Dunford et al., 2012). To date there has been no research into salt levels in fast foods in Costa Rica, despite research indicating that the contribution of these foods to the diet is higher than recommendations (Ministerio de Salud, 2011). Hence the aim of this study was to examine the reported salt content in products from popular fast food chains in Costa Rica, and to report findings by food category and by fast food chain.

## Materials and methods

### Data collection

This study comprised a survey of the salt and energy content of fast food menu items from fast food companies in Costa Rica, with data collection done in January 2013. Data were obtained from company websites. The 11 fast food companies most visited by Costa Rican families were identified: Burger King, Domino's Pizza, Kentucky Fried Chicken (KFC), McDonald's, Pizza Hut, Popeye's, Quiznos, Subway, Taco Bell, Teriyaki and Wendy's (Euromonitor International, 2011; Venegas & Villalobos, 2012). Four of these restaurants were excluded from analysis (Burger King, McDonald's, Quiznos and Wendy's) as they did not provide product serving size information and/or nutrition information per 100 g on the company website.

Data for 311 fast food items were collected and entered into a Microsoft Excel spreadsheet. For each product the following information was recorded: company name, product name, serving size (g), energy (kcal and kJ) and salt (g) content. Where information was available, we recorded energy and salt content both per serving and per 100 g. When both measures were not provided, one was calculated from the other using the serving size of the item. If sodium content was provided instead of salt, the value was converted by multiplying by 2.5 (Institute of Medicine, 2004). The data entry process was checked by selecting a random sample of 5% of entries and comparing the information in our database against the original website source.

### Definitions of product categories

Ten fast food categories were defined based on those used for previous report from the US (O'Donnell et al., 2008) and Australia (Dunford, 2012) and the product groupings used by the fast food industry: "chicken products" (nuggets, drumsticks, fried, grilles or roasted chicken); "pizza", "potato products" (smashed, fried, boiled with or without condiments); "rice, pasta and noodles", "salads" (all salad items and salads with additional ingredients); "sandwiches" (all kinds, wraps and burritos), "sauces" (all sauces, salad dressings and condiments), "tacos" (including nachos and chalupas), "dessert" (all types) and "others". The overarching goal for the categorization system was that it be applicable to industry and reflect consumer purchasing patterns. Breakfast meals were excluded from this analysis, because not all restaurants offer breakfast menus.

### Data analysis

Mean levels (and ranges) for energy and salt content per 100 g were calculated for each food category, overall and for each company.

We tested homogeneity of variances (Levene's test,  $p < 0.05$ ) to obtain a single  $p$ -value that indicated the presence or absence of variability in energy and salt content across food categories and fast food chains. Due to the variance observed, we used the Welch ANOVA for comparison of means of unequal variances ( $p < 0.05$ ). We used one-way analysis of variance to compare salt content between companies, using Tukey–Kramer test HSD. A  $p$  value  $\leq 0.05$  was considered statistically significant. All statistical analyses were performed using SPSS version 20.

## Results

Only one company, Taco Bell, out of the seven that were analyzed provided regional nutritional information in their websites. The others showed nutritional information from US, Canada or Europe. Salt and energy content data were collected for 311 fast food items from seven companies (Table 1). The number of products per company ranged from 12 (Domino's Pizza) to 75 (Taco Bell). There was a statistically significant correlation ( $p < 0.05$ ) between salt and energy content; with sodium content increasing with an increase in energy content.

Salt content varied significantly between companies ( $p < 0.05$ ) with Subway having the lowest mean level of salt (0.97 g/100 g) and Popeye's having the highest (1.57 g/100 g). There were significant differences ( $p < 0.05$ ) in salt content between Subway 0.97 g and, Pizza Hut 1.52 g, Popeye's 1.57 g and KFC 1.57 g/100 g (Fig. 1).

Figure 2 shows substantial variations in mean salt content were also found between food categories. For example, there was a more than fourfold difference in mean salt content per 100 g between "salads" (0.45 g) and "sauces" (2.16 g). A wide and significant ( $p < 0.05$ ) variation per 100 g was also seen within food categories. For example, salt content in "chicken" ranged from <1 g/100 g to 2.7 g/100 g and "sandwiches, wraps and burritos" ranged from 0.5 g/100 g to 2.1 g/100 g (Fig. 2).

## Discussion

The main finding from this study was the marked variability in the reported salt content of products by major fast food chains in Costa Rica. This was true for comparisons between companies, between different products and within food categories. However, the large range of salt content in products from the same food category shows that there is a clear opportunity for reformulation of products toward the lower end of the range by fast food companies to improve the healthiness of their offerings. For example, there was an almost threefold difference in salt content across chicken products and fivefold difference across sauces and condiments. This is in line with research from other countries such as Australia, the US, New Zealand, Canada, France and the UK (Dunford et al., 2012; Johnson et al., 2010), however this study represents the first one of its kind in Latin America (using this methodology) to evaluate

**Table 1**  
Energy (kcal) and salt (g) content (mean and range) per 100 g by company.

Companies	Products (N)	Energy (kcal)		Salt (g)	
		Mean	(SD)	Mean	(SD)
Domino's Pizza	12	265	(49.7)	1.33	(0.40)
KFC	50	236	(118.4)	1.57	(0.84)
Pizza Hut	59	260	(91.6)	1.52	(0.58)
Popeye's	37	206	(127.8)	1.57	(0.77)
Subway	49	213	(160.1)	0.97	(0.60)
Taco Bell	75	132	(89.0)	1.29	(0.36)
Teriyaki	29	225	(64.6)	1.36	(1.33)

Note: SD: Standard deviation. ANOVA test was used for comparisons of mean nutrient content between restaurants. In all cases  $p$  values were  $< 0.05$ .

Download English Version:

<https://daneshyari.com/en/article/7309689>

Download Persian Version:

<https://daneshyari.com/article/7309689>

[Daneshyari.com](https://daneshyari.com)