Accepted Manuscript



Importance of human right inferior frontoparietal network connected by inferior branch of superior longitudinal fasciculus tract in corporeal awareness of kinesthetic illusory movement

Kaoru Amemiya, Eiichi Naito, PhD

PII: S0010-9452(16)30011-9

DOI: 10.1016/j.cortex.2016.01.017

Reference: CORTEX 1687

To appear in: *Cortex*

Received Date: 26 May 2015

Revised Date: 14 December 2015

Accepted Date: 18 January 2016

Please cite this article as: Amemiya K, Naito E, Importance of human right inferior frontoparietal network connected by inferior branch of superior longitudinal fasciculus tract in corporeal awareness of kinesthetic illusory movement, *CORTEX* (2016), doi: 10.1016/j.cortex.2016.01.017.

Importance of human right inferior frontoparietal network connected by inferior branch of superior longitudinal fasciculus tract in corporeal awareness of kinesthetic illusory movement

Kaoru Amemiya[1,2,3] and Eiichi Naito[3,4]

[1]The Japan Society for the Promotion of Science, 5-3-1, Koujimachi, Chiyoda, Tokyo, 102-0083, Japan;

[2]Department of Biosciences and Informatics, Faculty of Science and Technology, Keio University, 3-14-1, Hiyoshi, Kohoku-ku, Yokohama, Kanagawa, 223-8522, Japan;

[3]Center for Information and Neural Networks (CiNet), National Institute of Information and Communications Technology (NICT), 2A6, 1-4 Yamadaoka, Suita, Osaka, 565-0871, Japan;

[4]Graduate School of Medicine & Graduate School of Frontier Biosciences, Osaka University, Suita, Osaka, Japan

Correspondence:
Eiichi Naito, PhD
Center for Information and Neural Networks (CiNet), National Institute of Information and Communications Technology (NICT), 2A6, 1-4 Yamadaoka, Suita, Osaka, 565-0871, Japan
Tel: +81-80-9098-3256
Fax: +81-6-7174-8612
Email: eiichi.naito@nict.go.jp