



Original Article

Reprint of Hormonal correlates of pathogen disgust: testing the compensatory prophylaxis hypothesis^{☆, ☆ ☆, ☆}

Benedict C. Jones^{a,*}, Amanda C. Hahn^b, Claire I. Fisher^a, Hongyi Wang^c, Michal Kandrik^{d,e}, Anthony J. Lee^a, Joshua M. Tybur^{d,e}, Lisa M. DeBruine^a

^a Institute of Neuroscience & Psychology, University of Glasgow, UK

^b Department of Psychology, Humboldt State University, USA

^c School of Psychology and Cognitive Science, East China Normal University, China

^d Department of Experimental and Applied Psychology, VU Amsterdam, Netherlands

^e Institute Brain and Behavior Amsterdam, Netherlands

A B S T R A C T

Raised progesterone during the menstrual cycle is associated with suppressed physiological immune responses, reducing the probability that the immune system will compromise the blastocyst's development. The Compensatory Prophylaxis Hypothesis proposes that this progesterone-linked immunosuppression triggers increased disgust responses to pathogen cues, compensating for the reduction in physiological immune responses by minimizing contact with pathogens. Although a popular and influential hypothesis, there is no direct, within-woman evidence for correlated changes in progesterone and pathogen disgust. To address this issue, we used a longitudinal design to test for correlated changes in salivary progesterone and pathogen disgust (measured using the pathogen disgust subscale of the Three Domain Disgust Scale) in a large sample of women ($N = 375$). Our analyses showed no evidence that pathogen disgust tracked changes in progesterone, estradiol, testosterone, or cortisol. Thus, our results provide no support for the Compensatory Prophylaxis Hypothesis of variation in pathogen disgust.

1. Introduction

Suppressed physiological immune responses during the luteal phase of the menstrual cycle (when raised progesterone prepares the body for pregnancy) reduce the probability that the immune system will compromise the development of the blastocyst (reviewed in Fleischman & Fessler, 2011). The Compensatory Prophylaxis Hypothesis proposes that this progesterone-linked immunosuppression is associated with increased disgust toward pathogen cues, compensating for the reduction in physiological immune responses by reducing the probability of contact with pathogens (Fessler, Eng, & Navarrete, 2005; Fleischman & Fessler, 2011; Żelaźniewicz, Borkowska, Nowak, & Pawłowski, 2016).

Fleischman and Fessler (2011) and Żelaźniewicz et al. (2016) have presented the strongest evidence for (and most direct tests of) the

Compensatory Prophylaxis Hypothesis to date. In both studies, women with higher progesterone levels reported stronger disgust toward pathogen cues. Another study reporting stronger disgust responses to pathogen cues during the first (i.e., highest-progesterone) trimester of pregnancy has also been interpreted as supporting the Compensatory Prophylaxis Hypothesis (Fessler et al., 2005). However, these three studies employed between-subject designs, which have been shown to be weak (e.g., underpowered) tests for hypotheses concerning hormone-linked changes in behavior (Gangestad et al., 2016) and allow only indirect tests of the hypothesis that within-woman changes in pathogen disgust and progesterone are correlated. The two studies that measured progesterone levels (Fleischman & Fessler, 2011; Żelaźniewicz et al., 2016) employed relatively small sample sizes (N s of 79 and 30, respectively), meaning that they were likely underpowered

DOI of original article: <http://dx.doi.org/10.1016/j.evolhumbehav.2017.12.004>

^{*} This article is a reprint of a previously published article. For citation purposes, please use the original publication details; Evolution and Human Behavior Volume 39, Issue 2, March 2018, Pages 166–169.

^{☆☆} This research was supported by ERC Grants awarded to BCJ (OCMATE), LMD (KINSHIP), and JMT (HBIS). The authors thank Lars Penke, Julia Jünger, and two anonymous reviewers for helpful comments on our manuscript.

^{*} BCJ, LMD, ACH, and CIF designed the study; ACH, CIF, HW, and MK collected data; LMD, BCJ, HW, MK, and AJL analyzed the data; BCJ, LMD, and JT drafted the manuscript; and all authors revised the manuscript.

^{*} Corresponding author.

E-mail address: ben.jones@glasgow.ac.uk (B.C. Jones).

<https://doi.org/10.1016/j.evolhumbehav.2018.05.006>

Received 11 October 2017; Accepted 12 December 2017

1090-5138/ Crown Copyright © 2017 Published by Elsevier Inc. All rights reserved.

(see Gangestad et al., 2016).

Other studies often cited as evidence for the Compensatory Prophylaxis Hypothesis are also problematic. For example, greater hostility to out-group individuals during the first trimester of pregnancy has been interpreted as evidence for the Compensatory Prophylaxis Hypothesis because out-group individuals putatively pose a greater pathogen threat than do in-group individuals (Navarrete, Fessler, & Eng, 2007). However, the hypothesis that hostility to out-group individuals necessarily reflects pathogen avoidance has recently been extensively critiqued (Aarøe, Osmundsen, & Petersen, 2016; de Barra & Curtis, 2012; Tybur et al., 2016). Reports that women show stronger aversions to individuals displaying facial cues of illness (e.g., pallor) at high-progesterone points in the menstrual cycle (Jones et al., 2005) have also been interpreted as evidence for the Compensatory Prophylaxis Hypothesis. These results were not replicated in a higher-powered study that directly tested for correlated changes in measured progesterone and aversions to facial cues of illness (Jones et al., 2017b).

In summary, considering its influence in both the emotion and endocrinology literatures, the Compensatory Prophylaxis Hypothesis is supported by a surprisingly weak body of evidence. In the current study, we rigorously tested the Compensatory Prophylaxis Hypothesis by using a longitudinal design to investigate whether within-woman changes in steroid hormone levels (including progesterone) and changes in components of disgust sensitivity (including pathogen disgust) were correlated in a large sample of women ($N = 375$). We assessed disgust sensitivity using Tybur, Lieberman, and Griskevicius's (2009) Three Domain Disgust Scale, which assesses disgust sensitivity in three different domains: pathogen disgust, sexual disgust, and moral disgust. The Compensatory Prophylaxis Hypothesis predicts that pathogen disgust will track changes in women's progesterone levels (Fessler et al., 2005; Fleischman & Fessler, 2011; Żelaźniewicz et al., 2016). Indeed, the studies testing the Compensatory Prophylaxis Hypothesis have each used either the Three Domain Disgust Scale, similar self-report measures of disgust or contamination sensitivity, or disgust ratings of images portraying cues to pathogens (Fessler et al., 2005; Fleischman & Fessler, 2011; Żelaźniewicz et al., 2016).

2. Methods

2.1. Participants

We tested 375 heterosexual women (mean age = 21.6 years, $SD = 3.3$ years), all of whom reported that they were not using any form of hormonal contraceptive (i.e., reported having natural menstrual cycles). Participants completed up to three blocks of test sessions. Each of the three blocks of test sessions consisted of five weekly test sessions. Women participated as part of a large study of possible effects of steroid hormones on women's behavior (Jones et al., 2017b). The data analyzed here are all responses from blocks of test sessions where women were not using any form of hormonal contraceptive and test sessions where they completed at least one subscale of Tybur et al.'s (2009) Three Domain Disgust Scale. Following these restrictions, 337 women had completed five or more test sessions and 98 of these women completed ten test sessions. Thirty-eight women completed fewer than five test sessions.

2.2. Three Domain Disgust Scale

Participants completed Tybur et al.'s (2009) Three Domain Disgust Scale in each test session. This 21-item measure asks participants to rate each of 21 actions from not at all disgusting (0) to extremely disgusting (6). The actions were divided into three domains: pathogen disgust (e.g., stepping on dog poop), sexual disgust (e.g., hearing two strangers having sex), and moral disgust (e.g., deceiving a friend). Question order was fully randomized. The full instructions for the questionnaire were: "The following items describe a variety of concepts. Please rate how

disgusting you find the concepts described in the items, where 0 means that you do not find the concept disgusting at all, and 6 means that you find the concept extremely disgusting."

The mean score on the pathogen disgust subscale was 25.99 ($SD = 7.98$), the mean score on the sexual disgust subscale was 19.95 ($SD = 8.71$), and the mean score on the moral disgust subscale was 27.82 ($SD = 8.32$). Intra-class correlation coefficients were high for each subscale (pathogen: 0.82; 95% CIs: 0.80, 0.85; sexual: 0.88; 95% CIs: 0.86, 0.89; moral = 0.79; 95% CIs: 0.76, 0.82). Consistent with past research (Olatunji et al., 2012), these intra-class correlation coefficients indicate that scores on the Three Domain Disgust Scale are stable over time (or, at least, over the time span sampled in the current study). Nevertheless, small fluctuations in disgust sensitivity could covary with variation in hormones.

2.3. Saliva samples

Participants provided a saliva sample via passive drool (Papacosta & Nassis, 2011) in each test session. Participants were instructed to avoid consuming alcohol and coffee in the 12 h prior to participation and avoid eating, smoking, drinking, chewing gum, or brushing their teeth in the 60 min prior to participation. Each woman's test sessions took place at approximately the same time of day to minimize effects of diurnal changes in hormone levels (Bao et al., 2003; Veldhuis et al., 1988).

Saliva samples were frozen immediately and stored at -32°C until being shipped, on dry ice, to the Salimetrics Lab (Suffolk, UK) for analysis, where they were assayed using the Salivary 17β -Estradiol Enzyme Immunoassay Kit 1-3702 ($M = 3.30$ pg/mL, $SD = 1.27$ pg/mL, sensitivity = 0.1 pg/mL, intra-assay CV = 7.13%, inter-assay CV = 7.45%), Salivary Progesterone Enzyme Immunoassay Kit 1-1502 ($M = 148.59$ pg/mL, $SD = 96.20$ pg/mL, sensitivity = 5 pg/mL, intra-assay CV = 6.20%, inter-assay CV = 7.55%), Salivary Testosterone Enzyme Immunoassay Kit 1-2402 ($M = 87.57$ pg/mL, $SD = 27.19$ pg/mL, sensitivity < 1.0 pg/mL, intra-assay CV = 4.60%, inter-assay CV = 9.83%), and Salivary Cortisol Enzyme Immunoassay Kit 1-3002 ($M = 0.23$ $\mu\text{g}/\text{dL}$, $SD = 0.16$ $\mu\text{g}/\text{dL}$, sensitivity < 0.003 $\mu\text{g}/\text{dL}$, intra-assay CV = 3.50%, inter-assay CV = 5.08%). Although Fleischman and Fessler (2011) and Żelaźniewicz et al. (2016) only reported progesterone in their studies, we measured and report analyses of estradiol, cortisol, and testosterone, in addition to progesterone, to test whether links between pathogen disgust and hormonal status are driven specifically by progesterone, as the Compensatory Prophylaxis Hypothesis proposes. Mean minimum and maximum hormone levels are given in our Supplemental information.

Hormone levels more than three standard deviations from the sample mean for that hormone or where Salimetrics indicated levels were outside the sensitivity range of their relevant ELISA were excluded from the dataset (~1% of hormone measures were excluded for these reasons). The descriptive statistics given above do not include these excluded values. Values for each hormone were centered on their subject-specific means to isolate effects of within-woman changes in hormones. They were then scaled (i.e., divided by a constant) so the majority of the distribution for each hormone varied from -0.5 to 0.5 to facilitate calculations in the linear mixed models. Since hormone levels were centered on their subject-specific means, women with only one value for a hormone could not be included in the analyses.

2.4. Analyses

Linear mixed models were used to test for possible effects of hormonal status on disgust sensitivity. Analyses were conducted using R version 3.3.2 (R Core Team, 2016), with lme4 version 1.1-13 (Bates, Maechler, Bolker, & Walker, 2014) and lmerTest version 2.0-33 (Kuznetsova, Brockhoff, & Christensen, 2013). The dependent variable was Three Domain Disgust subscale score (separate models were run for

Download English Version:

<https://daneshyari.com/en/article/7316206>

Download Persian Version:

<https://daneshyari.com/article/7316206>

[Daneshyari.com](https://daneshyari.com)