

Motor cognition in patients treated with subthalamic nucleus deep brain stimulation: Limits of compensatory overactivity in Parkinson's disease

Klara Hagelweide, Anna Schönberger, Lutz Kracht, Theo Gründler, Gereon R. Fink, Ricarda I. Schubotz



PII: S0028-3932(18)30307-5
DOI: <https://doi.org/10.1016/j.neuropsychologia.2018.07.007>
Reference: NSY6843

To appear in: *Neuropsychologia*

Received date: 31 October 2017
Revised date: 7 May 2018
Accepted date: 6 July 2018

Cite this article as: Klara Hagelweide, Anna Schönberger, Lutz Kracht, Theo Gründler, Gereon R. Fink and Ricarda I. Schubotz, Motor cognition in patients treated with subthalamic nucleus deep brain stimulation: Limits of compensatory overactivity in Parkinson's disease, *Neuropsychologia*, <https://doi.org/10.1016/j.neuropsychologia.2018.07.007>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Motor cognition in patients treated with subthalamic nucleus deep brain stimulation: Limits of compensatory overactivity in Parkinson's disease

Klara Hagelweide^{a,b,*}, Anna Schönberger^a, Lutz Kracht^c, Theo Gründler^{d,e}, Gereon R. Fink^{a,f}, Ricarda I. Schubotz^{a,b}

^aDepartment of Neurology, University Hospital Cologne, Kerpener Str. 62 D-50937 Cologne, Germany

^bDepartment of Psychology, University of Muenster, Fliednerstr. 21 D-48149 Münster, Germany

^cMax Planck Institute for Metabolism Research, Gleueler Str. 50 D-50931 Cologne, Germany

^dCenter for Behavioral Brain Sciences, Otto-von-Guericke-University Magdeburg, Universitätsplatz 2, 39106 Magdeburg, Germany

^eFaculty of Economics and Management, Otto-von-Guericke-University Magdeburg, Universitätsplatz 2, 39106 Magdeburg, Germany

^fInstitute of Neuroscience and Medicine (INM3), Cognitive Neuroscience, Research Centre Jülich, Leo-Brandt-Straße D-52425 Jülich, Germany

*Corresponding author. Klara Hagelweide Department of Psychology, University of Münster, Fliednerstr. 21, 48149 Münster, Germany. Tel.: + 49 251/83 34134; fax: +49 251/83 34104. klara.hagelweide@wwu.de

Abstract

Recent fMRI findings revealed that impairment in a serial prediction task in patients suffering from Parkinson's disease (PD) results from hypoactivity of the **SMA**. Furthermore, hyperactivity of the lateral premotor cortex sustained performance after withdrawal of medication. To further explore these findings, we here examined the impact of deep brain

Download English Version:

<https://daneshyari.com/en/article/7317498>

Download Persian Version:

<https://daneshyari.com/article/7317498>

[Daneshyari.com](https://daneshyari.com)