
Author's Accepted Manuscript

Sex modulate neurophysiological hormones correlates of visual temporal attention

Cornelia Kranczioch, Anja Lindig, Markus Hausmann



www.elsevier.com/locate/neuropsychologia

PII: S0028-3932(16)30286-X

DOI: http://dx.doi.org/10.1016/j.neuropsychologia.2016.08.004

Reference: NSY6095

To appear in: Neuropsychologia

Received date: 21 January 2016 Revised date: 1 August 2016 Accepted date: 4 August 2016

Cite this article as: Cornelia Kranczioch, Anja Lindig and Markus Hausmann Sex hormones modulate neurophysiological correlates of visual tempora , Neuropsychologia t n t n http://dx.doi.org/10.1016/j.neuropsychologia.2016.08.004

This is a PDF file of an unedited manuscript that has been accepted fo publication. As a service to our customers we are providing this early version o the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain

ACCEPTED MANUSCRIPT

Sex hormones modulate neurophysiological correlates of visual temporal attention

Cornelia Kranczioch^{a,b,*}, Anja Lindig^{a,1}, Markus Hausmann^c

^aSchool of Medicine and Health Sciences, Department of Psychology,

Neuropsychology Lab, University of Oldenburg, 26111 Oldenburg, Germany

^bResearch Center Neurosensory Science, University of Oldenburg, 26111

Oldenburg, Germany

^cDepartment of Psychology, Durham University, South Road, Durham DH1 3LE,

United Kingdom

cornelia.kranczioch@uni-oldenburg.de

anja.lindig@pius-hospital.de

markus.hausmann@durham.ac.uk

*Corresponding author

Abstract

The functional cerebral asymmetry (FCA) in processing targets within rapid serial visual presentation (RSVP) streams has been reported to fluctuate across the menstrual cycle, with identification of the second of two closely spaced targets being impaired when both targets occur in the left or the right hemifield stream during the luteal phase, while during the menstrual phase identification of the second target is only impaired for target pairs presented in

¹ Present address: Pius Hospital Oldenburg, Klinik für Hämatologie und Onkologie, Georgstraße 12, 26121 Oldenburg

Download English Version:

https://daneshyari.com/en/article/7318498

Download Persian Version:

https://daneshyari.com/article/7318498

<u>Daneshyari.com</u>