

## Accepted Manuscript

Mass segregation phenomena using the Hamiltonian Mean Field model

J.R. Steiner, , T.O. Zolacir, jr.

PII: S0378-4371(17)30988-3  
DOI: <https://doi.org/10.1016/j.physa.2017.09.095>  
Reference: PHYSA 18700

To appear in: *Physica A*

Received date: 20 March 2017  
Revised date: 22 September 2017

Please cite this article as: J.R. Steiner, T.O. Zolacir, Mass segregation phenomena using the Hamiltonian Mean Field model, *Physica A* (2017), <https://doi.org/10.1016/j.physa.2017.09.095>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Mass Segregation Phenomena using the Hamiltonian Mean Field Model

J.R. Steiner and Zolacir T. O. Jr

Highlights:

- The usage of Hamiltonian Mean Field model to the study of mass segregation phenomena.
- Show that the time scale for the QSS in systems with different masses is  $N$  and not  $N^2$  as for identical particles.
- We show the explicit emergence of mass segregation in one-dimensional system.

Download English Version:

<https://daneshyari.com/en/article/7376380>

Download Persian Version:

<https://daneshyari.com/article/7376380>

[Daneshyari.com](https://daneshyari.com)